Table S2. Stratified analysis of the association between nursing diagnoses and mortality according to the Shock Index

Nursing Diagnosis	Shock Index < 0.7		Shock Index ≥ 0.7		P for interaction
	Adjusted RR (95% CI)	p-value	Adjusted RR (95% CI)	p-value	
Risk for unstable blood glucose	0.15 (0.02, 1.10)	0.060	0.60 (0.19, 1.88)	0.388	0.001
Risk for electrolyte imbalance	2.32 (1.21, 4.45)	0.010	2.86 (1.39, 6.53)	0.005	0.004
Excess fluid volume	1.87 (0.77, 4.40)	0.169	2.88 (0.86, 9.59)	0.087	<0.001
Impaired gas exchange	2.04 (1.06, 3.92)	0.031	2.86 (1.48, 6.26)	0.002	<0.001
Decreased cardiac output	1.67 (1.21, 2.29)	0.002	1.85 (1.25, 2.70)	0.021	0.005
Risk for ineffective cerebral tissue perfusion	1.32 (0.76, 1.49)	0.283	7.31 (3.02, 17.68)	0.001	<0.001
Risk for impaired cardiovascular function	1.47 (0.84, 2.58)	0.171	2.35 (1.32, 4.17)	0.004	0.003
Anxiety	0.21 (0.03, 1.40)	0.109	0.58 (0.30, 1.42)	0.117	0.001
Risk for shock	3.09 (1.75, 5.49)	0.001	4.55 (1.77, 11.72)	0.002	<0.001
Acute pain	0.62 (0.25, 1.22)	0.297	0.16 (0.03, 0.79)	0.024	0.002
Impaired psychological comfort	0.29 (0.12, 0.67)	0.004	0.80 (0.10, 1.02)	0.052	0.001

Abbreviations: RR, risk ratio; CI, confidence interval.