

Supplementary Materials

Antioxidant Quercetin 3-O-Glycosylated Plant Flavonols Contribute to Transthyretin Stabilization

Lidia Ciccone ^{1,*}, Nicolò Tonali ², Carole Fruchart-Gaillard ³, Lucia Barlettani ¹, Armando Rossello ^{1,4},
Alessandra Braca ^{1,5}, Elisabetta Orlandini ^{4,6} and Susanna Nencetti ^{1,5}

¹ Department of Pharmacy, University of Pisa, 56126 Pisa, Italy; l.barlettani@studenti.unipi.it (L.B.), armando.rossello@unipi.it (A.R.), alessandra.braca@unipi.it (A.B.), susanna.memcetti@unipi.it (S.N.)

² CNRS, BioCIS, Université Paris-Saclay, 92290 Châtenay-Malabry, France; nicolo.tonali@universite-paris-saclay.fr

³ Département Médicaments et Technologies pour la Santé (DMTS), SIMoS, CEA, INRAE, Université Paris Saclay, 91191 Gif-Sur-Yvette, France, carole.fruchart@cea.fr

⁴ Research Center "E. Piaggio", Università di Pisa, 56122 Pisa, Italy, elisabetta.orlandini@unipi.it

⁵ Interdepartmental Research Centre Nutraceuticals and Food for Health (NUTRAFOOD), University of Pisa, 56126 Pisa, Italy

⁶ Department of Earth Sciences, University of Pisa, 56100 Pisa, Italy

* Correspondence: lidia.ciccone@unipi.it

Table of Contents

Figure S1. OMIT map of QUE 3-O- β -D-galactopyranoside




Figure S1. Screenshot of COOT graphic interface. OMIT map of QUE 3-O- β -D-galactopyranoside contoured at 3σ .