

Camphor Sulfonylimine Compounds: Bottom-Up

Design of MOF's from Organic Frameworks Based on X-rayS and DFT-D3

Joana P. Costa ^a, M. Fernanda N.N. Carvalho ^a and Adelino M. Galvão ^{b,*}

^a Centro de Química Estrutural, Institute of Molecular Sciences and Departamento de Engenharia Química, Instituto Superior Técnico, Universidade de Lisboa, Avenida António José de Almeida, n.º 12, 1000-043 Lisboa, Portugal.

^b Centro de Química Estrutural, Institute of Molecular Sciences and Departamento de Engenharia Química, Instituto Superior Técnico, Campus Taguspark, Universidade de Lisboa, Av. Prof. Doutor Cavaco Silva 2744-016 Porto Salvo, Portugal,

* Correspondence: E-mail: adelino@tecnico.ulisboa.pt.

Supplementary data

Spectroscopic information: ¹H, ¹³C NMR and FTIR experimental data is provided for coordination compounds C1 and C2.

Coordination compound C1




Fig. S1 – NMR ¹H spectrum of compound C1.




Fig. S2 – NMR ¹³C spectrum of compound C1.




Fig. S3 – FTIR spectrum of compound C1.

Coordination compound **C2**




Fig. S4 – NMR ¹H spectrum of compound C2.




Fig. S5 – NMR ¹³C spectrum of compound C2.




Fig. S6 – FTIR spectrum of compound C2.