SUPPLEMENTARY MATERIAL

Log, rock, and a gradient of choice: *Speleomantes strinatii* adjusts shelter use with stream distance

Giacomo Rosa[1,2,*], Andrea Costa[1], Sebastiano Salvidio[1]

[1] Department of Earth, Environment and Life Sciences (DISTAV), University of Genova, Corso Europa 26, 16132 Genova, Italy; andrea-costa-@hotmail.it (A.C.); sebastiano.salvidio@unige.it (S.S.)

[2] National Biodiversity Future Centre (NBFC), Palermo, Italy

* Correspondence: giacomorosa@live.it; giacomo.rosa@edu.unige.it

Table S1 Summary table regarding the available refuges and their use in the three forest habitats located at different distances from the stream, provided in meters (m) between brackets.

	Availability			Usage occasions		
	Chestnut (2-15 m)	Transitional (22-42 m)	Mixed (45-67 m)	Chestnut (2-15 m)	Transitional (22-42 m)	Mixed (45-67 m)
Trunk	15	9	6	20	6	2
Rock	2	4	35	1	1	23
Total	17	13	41	21	7	25



Figure S1 Relationship between mean number of available woody shelters and distance from the watercourse.



Figure S2 Distribution of shelters' surface coverage for logs (a) and rocks (b) respectively. Bars represent 95% confidence intervals.