

# **In Comparison to Pathological Q Waves, Selvester Score is a Superior Diagnostic Indicator of Increased Long-Term Mortality Risk in ST Elevation Myocardial Infarction Patients Treated with Primary Coronary Intervention**

**Maria Holicka <sup>1,2</sup>, Pavla Cuckova <sup>1,2</sup>, Katerina Hnatkova <sup>3</sup>, Lumir Koc <sup>1,2</sup>, Tomas Ondrus <sup>1,2</sup>, Petr Lokaj <sup>1,2</sup>, Jiri Parenica <sup>1,2</sup>, Tomas Novotny <sup>1,2,\*</sup>, Petr Kala <sup>1,2</sup>, and Marek Malik <sup>2,3</sup>**

<sup>1</sup> Department of Internal Medicine and Cardiology, University Hospital Brno, Jihlavská 20, 625 00 Brno, Czech Republic; holicka.maria@fnbrno.cz (M.H.); cuckova.pavla@fnbrno.cz (P.C.); koc.lumir@fnbrno.cz (L.K.); ondrus.tomas@fnbrno.cz (T.O.); lokaj.petr2@fnbrno.cz (P.L.); parenica.jiri@fnbrno.cz (J.P.); kala.petr@fnbrno.cz (P.K.)

<sup>2</sup> Department of Internal Medicine and Cardiology, Faculty of Medicine, Masaryk University, Jihlavská 20, 625 00 Brno, Czech Republic; marek.malik@imperial.ac.uk

<sup>3</sup> National Heart and Lung Institute, Imperial College of London, 72 Du Cane Rd, Shepherd's Bush, London W12 0NN, UK; k.hnatkova@imperial.ac.uk

\* Correspondence: novotny.tomas3@fnbrno.cz; Tel.: +420-53223-2454

**Supplemental Table 1. QRS scoring system for estimating infarct size - the Selvester Score**  
(see Ref. 9)

Lead	Q duration (msec)	R duration (msec)	Ratios	Points	Lead max.	
I	≥30			1	2	
			R/Q≤1	1		
II	≥40			2	2	
	≥30			1		
aVL	≥30			1	2	
			R/Q≤1	1		
aVF	≥50			3	5	
	≥40			2		
	≥30			1		
			R/Q≤1	2		
			R/Q≤2	1		
V1	any			1	4	
		≥50				2
		≥40				1
				R/S≥1		1
V2	any	≤20		1	4	
		≥60				2
		≥50				1
				R/S≥1,5		1
V3	any	≤30		1	1	
V4	≥20			1	3	
			R/Q or R/S ≤0,5	2		
			R/Q or R/S ≤1,5	1		
V5	≥30		R/Q or R/S ≤1	1	3	
			R/Q or R/S ≤3	2		
				1		
V6	≥30			1	3	
			R/Q or R/S ≤1	2		
			R/Q or R/S ≤3	1		
Lead max – the maximum point count for particular lead						