

Supplementary Material

From euglycemia to recent onset of Type 2 diabetes mellitus: circulating microRNA profiling reveals distinct, and early microRNA signatures

Marta Greco, Maria Mirabelli, Alessandro Salatino, Francesca Accattato, Vincenzo Aiello, Francesco S. Brunetti, Eusebio Chiefari, Salvatore A. Pullano, Antonino S. Fiorillo, Daniela P. Foti and Antonio Brunetti

Table S1. Highly expressed miRNAs (Ct \leq 25) in euglycemic controls, and IFG and new-onset T2D patients.

Control		IFG		New-onset T2D	
Mature ID	Ct	Mature ID	Ct	Mature ID	Ct
<i>hsa-miR-328-3p</i>	22.58	<i>hsa-miR-451a</i>	22.04	<i>hsa-miR-451a</i>	21.68
<i>hsa-miR-486-5p</i>	24.46	<i>hsa-miR-486-5p</i>	23.46	<i>hsa-miR-486-5p</i>	22.36
<i>hsa-miR-92a-3p</i>	24.93	<i>hsa-miR-223-3p</i>	23.47	<i>hsa-miR-92a-3p</i>	23.20
		<i>hsa-miR-92a-3p</i>	24.27	<i>hsa-miR-223-3p</i>	23.35
				<i>hsa-miR-373-5p</i>	24.11
				<i>hsa-miR-1280</i>	24.32
				<i>hsa-miR-4454</i>	24.59