

Table S1. Differences in the patient interval (in days) based on sociodemographic and clinical factors.

Characteristic		<i>n</i>	Interval Median (IQR) (In Days)	<i>p</i> -Value ¹
Age	<60	49	63 (21–181)	0.975
	>60	51	61 (21–180)	
Gender	Male	39	60 (19–102)	0.157
	Female	61	90 (27–181)	
Race	Malay	21	66 (30–150)	0.690
	Chinese	19	60 (21–360)	
	Indian	50	76 (23–180)	
	Indigenous	10	42 (9–92)	
Location	Urban	46	60 (19–181)	0.566
	Rural	54	90 (21–180)	
Education	None/Primary	45	63 (30–181)	0.224
	Secondary/Tertiary	55	61 (18–120)	
Occupation	Not employed	43	90 (16–183)	0.409
	Employed/Retired	57	61 (27–102)	
Household income	≤MYR 4360	90	63 (21–181)	0.428
	>MYR 4360	10	57 (21–90)	
Anatomic site	Buccal mucosa	54	70 (19–181)	0.182
	Tongue	25	66 (33–120)	
	Others ²	21	31 (15–92)	
Diagnosis	OPMD	50	72 (19–120)	0.820
	Oral Cancer	47	61 (21–183)	

¹ Significant difference in patient interval between OPMD and oral cancer based on the Kruskal-Wallis H test with significance set to $p < 0.05$. ² Consists of the alveolar, gingiva, lip, floor of mouth, palate, mandible, and other sites.