

Eric F. Karlén

q = 1' based indices measure allelic differentiation, 'q = 2' based indices don't

Data Set I

Nine (9) pairs of artificial diploid populations (subsets) having ten samples per population, no allelic overlap between populations, and with equal AMD (D), equal heterozygosity, and equal variance within each population. By design, W_{AP} was 50% and the TDAP formatted indices were at the theoretical maximum for each subset.

N_a = total # alleles per subset
 N_{a-MAX} = theoretical maximum N_a per subset
 $N'_a = N_a$ expressed as a proportion of N_{a-MAX}

$N_{a-MAX} = 40$

Each data set is based on one 'marker'.

Alleles for each sample are p_i

		Data subsets																				
		SS-.9		SS-.8		SS-.7		SS-.6		SS-.5		SS-.4		SS-.3		SS-.2		SS-.1		SS-.05	Min SS	Max SS
		0.9		0.8		0.7		0.6		0.5		0.4		0.3		0.2		0.1		0.05	0.025	1
Sample	Population	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
1	A	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	11
2	A	1	12	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	12
3	A	3	13	1	13	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	13
4	A	4	14	4	14	1	14	1	14	1	1	1	1	1	1	1	1	1	1	1	1	14
5	A	5	15	5	15	5	15	1	15	1	1	1	1	1	1	1	1	1	1	1	1	15
6	A	6	16	6	16	6	16	6	16	1	16	1	1	1	1	1	1	1	1	1	1	16
7	A	7	17	7	17	7	17	7	17	7	17	1	17	1	1	1	1	1	1	1	1	17
8	A	8	18	8	18	8	18	8	18	8	18	8	18	1	18	1	1	1	1	1	1	18
9	A	9	19	9	19	9	19	9	19	9	19	9	19	9	19	1	19	1	1	1	1	19
10	A	10	20	10	20	10	20	10	20	10	20	10	20	10	20	10	20	1	20	1	20	20
11	B	51	51	51	51	51	51	51	51	51	51	51	51	51	51	51	51	51	51	51	51	61
12	B	51	62	51	51	51	51	51	51	51	51	51	51	51	51	51	51	51	51	51	51	62
13	B	53	63	51	63	51	51	51	51	51	51	51	51	51	51	51	51	51	51	51	51	63
14	B	54	64	54	64	51	64	51	64	51	51	51	51	51	51	51	51	51	51	51	51	64
15	B	55	65	55	65	55	65	51	65	51	51	51	51	51	51	51	51	51	51	51	51	65
16	B	56	66	56	66	56	66	56	66	51	66	51	51	51	51	51	51	51	51	51	51	66
17	B	57	67	57	67	57	67	57	67	57	67	51	67	51	51	51	51	51	51	51	51	67
18	B	58	68	58	68	58	68	58	68	58	68	58	68	51	68	51	51	51	51	51	51	68
19	B	59	69	59	69	59	69	59	69	59	69	59	69	59	69	51	69	51	51	51	51	69
20	B	60	70	60	70	60	70	60	70	60	70	60	70	60	70	60	70	51	70	51	70	70

Eric F. Karlin

q = 1' based indices measure allelic differentiation, 'q = 2' based indices don't

Data Set II

Ten (10) pairs of artificial populations (subsets) having the same parameters as the subsets in Data Set I.

with the exceptions of (1) allelic overlap occurring between the two populations in each subset and

(2) TDAP indices were not all at the theoretical maximum.

N_a = total # alleles per subset

N_{a-MAX} = theoretical maximum N_a per subset

N_{a-MAX} = 40

N'_a = N_a expressed as a proportion of N_{a-MAX}

Each data set is based on one 'marker'.

Alleles for each sample are placed in columns

Data subsets

		SS-93		SS-88		SS-78		SS-68		SS-58		SS-48		SS-38		SS-28		SS-18		SS-08	
		0.925		0.875		0.775		0.675		0.575		0.475		0.375		0.275		0.175		0.075	
Sample	Population																				
1	a	1	11	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
2	a	2	12	1	12	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
3	a	3	13	3	13	1	13	1	1	1	1	1	1	1	1	1	1	1	1	1	
4	a	4	14	4	14	4	14	1	14	1	1	1	1	1	1	1	1	1	1	1	
5	a	5	15	5	15	5	15	5	15	1	15	1	1	1	1	1	1	1	1	1	
6	a	6	16	6	16	6	16	6	16	6	16	1	16	1	1	1	1	1	1	1	
7	a	7	17	7	17	7	17	7	17	7	17	7	17	1	17	1	1	1	1	1	
8	a	8	18	8	18	8	18	8	18	8	18	8	18	8	18	1	18	1	1	1	
9	a	9	19	9	19	9	19	9	19	9	19	9	19	9	19	9	19	1	19	1	
10	a	10	20	10	20	10	20	10	20	10	20	10	20	10	20	10	20	10	20	1	
11	b	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
12	b	1	62	1	62	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
13	b	53	63	53	63	1	63	1	1	1	1	1	1	1	1	1	1	1	1	1	
14	b	54	64	54	64	54	64	1	64	1	1	1	1	1	1	1	1	1	1	1	
15	b	55	65	55	65	55	65	55	65	1	65	1	1	1	1	1	1	1	1	1	
16	b	56	66	56	66	56	66	56	66	56	66	1	66	1	1	1	1	1	1	1	
17	b	57	67	57	67	57	67	57	67	57	67	57	67	1	67	1	1	1	1	1	
18	b	58	68	58	68	58	68	58	68	58	68	58	68	58	68	1	68	1	1	1	
19	b	59	69	59	69	59	69	59	69	59	69	59	69	59	69	59	69	1	69	1	
20	b	60	70	60	70	60	70	60	70	60	70	60	70	60	70	60	70	60	70	1	

SS-05		Min SS		Max SS	
0.05		0.025		1	
1	1	1	1	1	11
1	1	1	1	1	12
1	1	1	1	1	13
1	1	1	1	1	14
1	1	1	1	1	15
1	1	1	1	1	16
1	1	1	1	1	17
1	1	1	1	1	18
1	1	1	1	1	19
1	1	1	1	1	20
51	51	1	1	51	61
51	51	1	1	52	62
51	51	1	1	53	63
51	51	1	1	54	64
51	51	1	1	55	65
51	51	1	1	56	66
51	51	1	1	57	67
51	51	1	1	58	68
51	51	1	1	59	69
51	51	1	1	60	70

Eric F. Karlén $q = 1'$ based indices measure allelic differentiation, ' $q = 2'$ based indices don't

Data Set III

Ten (10) pairs of artificial populations (subsets) were created based on the same parameters used for the subsets in DS-II except that heterozygosity, variance, and Δ are not balanced between the two populations in each subset.

N_s = total # alleles per subset
 $N_{s\text{-}MAX}$ = theoretical maximum N_s per subset
 $N'_s = N_s$ expressed as a proportion of $N_{s\text{-}MAX}$

Each data set is based on one 'marker'.
Alleles for each sample are placed in columns

		Data subsets																				0.025			
		SS-9		SS-8		SS-7		SS-6		SS-5		SS-4		SS-3		SS-2		SS-1		SS-05		Min SS		Max SS	
		0.9		0.8		0.7		0.6		0.5		0.4		0.3		0.2		0.1		0.05		0.025		1.0	
		N'_s																							
1	a	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
2	a	1	12	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
3	a	3	13	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
4	a	4	14	1	14	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
5	a	5	15	5	15	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
6	a	6	16	6	16	1	16	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
7	a	7	17	7	17	7	17	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
8	a	8	18	8	18	8	18	1	18	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
9	a	9	19	9	19	9	19	9	19	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
10	a	10	20	10	20	10	20	10	20	1	20	1	1	1	1	1	1	1	1	1	1	1	1	1	1
11	b	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
12	b	52	62	52	62	52	62	52	62	63	62	1	1	1	1	1	1	1	1	1	1	1	1	1	1
13	b	53	63	53	63	53	63	53	63	53	63	1	63	1	1	1	1	1	1	1	1	1	1	1	1
14	b	54	64	54	64	54	64	54	64	54	64	54	64	1	1	1	1	1	1	1	1	1	1	1	1
15	b	55	65	55	65	55	65	55	65	55	65	55	65	1	65	1	1	1	1	1	1	1	1	1	1
16	b	56	66	56	66	56	66	56	66	56	66	56	66	56	66	1	1	1	1	1	1	1	1	1	1
17	b	57	67	57	67	57	67	57	67	57	67	57	67	57	67	1	67	1	1	1	1	1	1	1	1
18	b	58	68	58	68	58	68	58	68	58	68	58	68	58	68	58	68	1	1	1	1	1	1	1	1
19	b	59	69	59	69	59	69	59	69	59	69	59	69	59	69	59	69	1	1	1	1	1	1	1	1
20	b	60	70	60	70	60	70	60	70	60	70	60	70	60	70	60	70	60	70	1	70	1	70	1	70