energies

MDPI

Article

Electrocatalytic Activity of Reduced Graphene Oxide Supported Cobalt Cinnamate for Oxygen Evolution Reaction

Myung Jun Lee †, Junyeop Kim †, Jaeun Kang, Hyewon Shin, Junghwan Do * and Seong Jung Kwon *

Department of Chemistry, Konkuk University, 120 Neungdong-ro Gwangjin-gu, Seoul 05029, Korea; hotbard@naver.com (M.J.L.); skykjyz@konkuk.ac.kr (J.K.); dolphinblue@naver.com (J.K.); shyewon9711@gmail.com (H.S.)

* Correspondence: junghwan@konkuk.ac.kr (J.D.); sjkwon@konkuk.ac.kr (S.J.K.); Tel.: +82-2-450-0429 (S.J.K)

† These authors are contributed equally.

Contents:

Figure S1: Transmission electron microscope (TEM) and energy dispersive spectroscopy (EDS) image of the cobalt cinnamate/rGO composite after electrochemical measurement.

Figure S2. TEM image of the cobalt cinnamate after electrochemical measurement.

Figure S3: The equivalent circuit for the calculation of EIS parameters.

Figure S4: XPS spectra for (a) C 1s, (b) O 1s, and (c) Co 2p regions using the catalyst/rGO composite-modified indium tin oxide electrode before and after electrochemical measurements.




Figure S1. Transmission electron microscope (TEM) and energy dispersive spectroscopy (EDS) image of the cobalt cinnamate/rGO composite after electrochemical measurement.




Figure S2. TEM image of the cobalt cinnamate after electrochemical measurement.




Figure S3. The equivalent circuit for the calculation of EIS parameters.




Figure S4. XPS spectra for (**a**) C 1s, (**b**) O 1s, and (**c**) Co 2p regions using the catalyst/rGO composite-modified indium tin oxide electrode before and after electrochemical measurements.