

Article

Impact of Polarization Reversal during Photoelectrocatalytic Treatment of WWTP Effluents

Maria Cristina Collivignarelli ^{1,2}, Marco Carnevale Miino ¹, Francesca Maria Caccamo ¹, Alessandro Abbà ^{3,*}, Massimiliano Bestetti ⁴, Silvia Franz ⁴

¹ Department of Civil Engineering and Architecture, University of Pavia, via Ferrata 3, 27100 Pavia, Italy; mcristina.collivignarelli@unipv.it (M.C.C.); marco.carnevalemiino01@universitadipavia.it (M.C.M.); francescamaria.caccamo01@universitadipavia.it (F.M.C.)

² Interdepartmental Centre for Water Research, University of Pavia, via Ferrata 3, 27100 Pavia, Italy

³ Department of Civil, Environmental, Architectural Engineering and Mathematics, University of Brescia, via Branze 43, 25123 Brescia, Italy

⁴ Department of Chemistry, Materials and Chemical Engineering "Giulio Natta", Politecnico di Milano, Via Mancinelli 7, 20131 Milano, Italy; massimiliano.bestetti@polimi.it (M.B.); silvia.franz@polimi.it (S.F.)

* Correspondence: alessandro.abba@unibs.it (A.A.)

FIGURE S1:




Figure S1. UV-Vis absorbance spectra of the untreated WWTP effluent.

FIGURE S2:




Figure S2. Emission spectra of medium-pressure Hg vapor lamp used in the study.

FIGURE S3:




Figure S3. Samples of untreated water and after 2h of EC, PL, PC, PEC and PECr.