Ecotoxicity of Fluoxetine Hydrochloride on Aquatic Organisms from Different Taxonomic Groups

Gabriel Souza-Silva 1,*, Maria Clara V. M. Starling 2, Clessius Ribeiro de Souza 1, Daniela de Carvalho 1, Mariângela Domingos Alcântara 1, Fernando Rodrigues-Silva 2, Cíntia Aparecida de Jesus Pereira 3, Marcos Paulo Gomes Mol 4 and Micheline Rosa Silveira 1

1 - Faculty of Pharmacy, Federal University of Minas Gerais, Belo Horizonte 31270-901, Brazil; clessiusdcq@gmail.com (C.R.d.S.); danielacarvalho@gmail.com (D.d.C.); mariangelaalcantara9@gmail.com (M.D.A.); michelinerosa@gmail.com (M.R.S.)

2 - Research Group on Environmental Applications of Advanced Oxidation Processes, Department of Sanitary and Environmental Engineering, Engineering School, Federal University of Minas Gerais, Belo Horizonte 31270-901, Brazil; mariaclara@desa.ufmg.br (M.C.V.M.S.); fernando.rodrigues002@gmail.com (F.R.d.S.)

3 - Department of Parasitology, Institute of Biological Sciences, Federal University of Minas Gerais, Belo Horizonte 31270-901, Brazil; cintiajp@icb.ufmg.br

4 - Department of Research and Development, Ezequiel Dias Foundation, Belo Horizonte 30510-080, Brazil; marcos.mol@funed.mg.gov.br

* Correspondence: silva_gs@yahoo.com

Table S1 - Physicochemical properties of fluoxetine.

CAS#	Molecular structure	Molecular weight	Formula	pKa	log Kow
54910-89-3		309.33 g/mol	$C_{17}H_{18}F_3NO$	9.8	4.1