Note: The images inside this file are for Figure 5 in the article below.

fermentation

MDPI

Article

Enhancement of biomethane yield from spent mushroom substrate: Biological pretreatment with the chlamydospores of Trichoderma viride

Wentao ZHU [a,1], Xianzhi LAI [a,1], Changfa LIU [a], Xiao WU [a], Xiaochen BAI [a], Yafan CAI [b], Xiaoling ZHAO [b], Zhe LI [a], Yongren HAO [a], Yanhua HUANG [a,*], Zehui ZHENG [a,*], Jie CHU [a]

[a] Biology Institute, Qilu University of Technology, Shandong Academy of Sciences, Jinan, 250103, China.
[b] School of Chemical Engineering, Zhengzhou University, Zhengzhou, 450001, China
* Correspondence: E-mail address: zhengzh@qlu.edu.cn (Z. Zheng); hyhua0827@126.com (Y. Huang)
[1] Wentao Zhu and Xianzhi Lai contributed equally to this work.




Figure. 5a The LEfSe evolutionary branches of fungal communities on the 1st day




Figure. 5b The LEfSe evolutionary branches of fungal communities on the 20th day




Figure. 5c The LEfSe evolutionary branches of bacterial communities on the 1st day



Figure. 5d The LEfSe evolutionary branches of bacterial communities on the 20th day




Figure. 5e The LEfSe evolutionary branches of archaeal communities on the 1st day




Figure. 5f The LEfSe evolutionary branches of archaeal communities on the 20th day