Supplementary materials

A CRISPR-Cas12a–based assay for efficient quantification of *Lactobacillus panis* in Chinese Baijiu brewing microbiome

Yanfeng Liu[1,3,4,5*], Mengchuang Wang[1,3,4,5], Fan Yang[2], Xiaolong Zhang[1,3,4,5], Jianghua Li[1,3,4,5], Guocheng Du[1,3,4,5], Li Wang[2,6*], and Jian Chen[3,4,5]

1. Key Laboratory of Carbohydrate Chemistry and Biotechnology, Ministry of Education, School of Biotechnology, Jiangnan University, Wuxi 214122, China

2. Science Center for Future Foods, Jiangnan University, Wuxi 214122, China

3. Engineering Research Center of Ministry of Education on Food Synthetic Biotechnology, Jiangnan University, Wuxi 214122, China

4. Jiangsu Province Engineering Research Center of Food Synthetic Biotechnology, Jiangnan University, Wuxi 214122, China

5. Kweichow Moutai Distillery Co., Ltd., Renhuai 564501, China

6. Kweichow Moutai Group, Renhuai 564501, China

*Corresponding author:

Yanfeng Liu

E-mail: yanfengliu@jiangnan.edu.cn

Li Wang

E-mail: wl@moutai.com.cn

Supplementary Figure:




Figure S1. Sequence region (coding vs. noncoding strand, 5’ vs 3’ ends) of the 16S rDNA that crRNA targets for enhanced detection.