

Supplementary materials

High-level production of catechol from glucose by engineered *Escherichia coli*

Guotian Song^{1,2,3}, Fengli Wu^{1,2}, Yanfeng Peng^{1,2}, Xiaolong Jiang^{1,2} and Qinhong Wang^{1,2,*}

¹CAS Key Laboratory of Systems Microbial Biotechnology, Tianjin Institute of Industrial Biotechnology, Chinese Academy of Sciences, Tianjin 300308, China; song_gt@tib.cas.cn (G.S.); wu_fl@tib.cas.cn (F.W.); peng_yf@tib.cas.cn (Y.P.); jiang_xl@tib.cas.cn (X.J.)

² National Center of Technology Innovation for Synthetic Biology, Tianjin 300308, China

³School of Life Science, University of Chinese Academy of Sciences, Beijing 100049, China

* Correspondence should be addressed to Qinhong Wang

E-mail:wang_qh@tib.cas.cn

Mailing address: Tianjin Institute of Industrial Biotechnology, Chinese Academy of Sciences,
32 XiQiDao, Tianjin Airport Economic Area, Tianjin 300308, P. R. China

Phone: +86-022-84861950

Supplementary Figures




Figure S1. The residual PCA after producing catechol via two-step whole-cell biocatalysis. Data are the average mean of triplicate samples, and the error bars represent standard deviations.




Figure S2. The mechanism of CA toxicity to microbial cells.