

## Article

# Anticholinesterase Activities of Different Solvent Extracts of Brewer's Spent Grain

Rares I. Birsan <sup>1,2</sup>, Peter Wilde <sup>2</sup>, Keith W. Waldron <sup>3</sup> and Dilip K. Rai <sup>1,\*</sup>

<sup>1</sup> Department of Food BioSciences, Teagasc Food Research Centre Ashtown, D15KN3K Dublin, Ireland; Rares.Birsan@teagasc.ie

<sup>2</sup> Food Innovation and Health Programme, Quadram Institute Bioscience, Norwich Research Park, Colney NR4 7UQ, UK; Pete.Wilde@quadram.ac.uk

<sup>3</sup> Anglia Science Writing Ltd., Wrappingham NR18 0RU, UK; keithwwaldron@outlook.com

\* Correspondence: dilip.rai@teagasc.ie; Tel.: +353-01-805-9500

## Supplementary Materials:

**Table S1.** Blends of individual polyphenols at 1 mg/mL mimicking their abundance in BSG fractions.

Sample	Total (μg/mL)	FA	p-CA	Cat	CafA	4-HBA	ProA
Blend FP EtOAc 1	1000	36.6	3.7	683.7	23.5	35.5	217
Blend BP DE 1	1000	633.4	354.5	-	7.1	4.5	0.5
Blend BP EtOAc 3	1000	703.2	267.6	-	19.9	5	4.3

FP—free phenolic extract, BP—bound phenolic extract followed by the fraction replicate number; μg/mL—microgram per milliliter; FA—ferulic acid, p-CA—coumaric acids, Cat—catechin, CafA—cafeic acid, 4-HBA—hydroxybenzoic acid, ProA—protocatechuic acid.

**Table S2.** Correlation coefficients among analyzed variables of BSG BP fractions.

Pearson Correl.	AChE	BChE	TPC	SQP	FA	p-CA	Cat	CafA	4-HBA	ProA	DeCa-DiFA	DiFA	TriFA
AChE	1												
BChE	0.687**	1											
TPC	0.375	0.511*	1										
SQP	0.543*	0.787**	0.904**	1									
FA	0.502*	0.736**	0.916**	0.990**	1								
p-CA	0.645**	0.869**	0.826**	0.980**	0.958**	1							
Cat	-0.036	-0.013	-0.293	-0.194	-0.189	-0.187	1						
CafA	0.163	0.428	0.922**	0.856**	0.887**	0.739**	-0.163	1					
4HBA	0.461*	0.715**	0.927**	0.985**	0.994**	0.941**	-0.190	0.907**	1				
ProA	-0.113	0.094	0.801**	0.610**	0.656**	0.439*	-0.139	0.922**	0.700**	1			
DeCa-DiFA	0.754**	0.896**	0.432	0.703**	0.621**	0.817**	-0.135	0.256	0.585**	-0.095	1		
DiFA	-0.229	-0.052	0.702**	0.481*	0.538*	0.298	-0.107	0.854**	0.585**	0.983**	-0.253	1	
TriFA	-0.227	-0.040	0.700**	0.488*	0.545*	0.306	-0.102	0.858**	0.595**	0.985**	-0.245	0.999**	1

Correlation is significant at the 0.01\*\* level and at the 0.05\* level (2-tailed).

**Table S3.** Summary of multiple regression model of AChE and BChE.

Model	R	R Square	Adjusted R Square	Std. Error of the Estimate
AChE	0.842 <sup>a</sup>	0.709	0.471	7.55749
AChE	0.763 <sup>b</sup>	0.582	0.536	7.08868
BChE	0.984 <sup>a</sup>	0.967	0.941	3.74240
BChE	0.933 <sup>b</sup>	0.871	0.856	5.82498

<sup>a</sup>Predictors: (Constant), FA, *p*-CA, Cat, CafA, 4HBA, ProA, DeCa-DiFA, TriFA, TPC; <sup>b</sup>Predictors: (Constant), FA, *p*-CA.

**Table S4.** The first four factor loadings for illustrating the interpretation of figure 2. Note: Numbers in bold represent the factor loading higher than 0.30.

Variable	PC1	PC2	PC3	PC4
FA	0.357	0.053	0.020	0.113
<i>p</i> -CA	0.340	0.176	-0.041	0.205
Cat	-0.049	-0.209	-0.765	0.004
CafA	0.343	-0.167	0.096	-0.028
4-HBA	0.356	-0.001	-0.068	0.111
ProA	0.163	-0.388	-0.488	-0.125
DeCa-DiFA	0.229	0.395	-0.164	0.362
DiFA	0.247	-0.383	0.226	-0.223
TriFA	0.248	-0.383	0.222	-0.224
ACHE	0.108	0.410	-0.087	-0.737
BCHE	0.223	0.339	-0.154	-0.344
TPC	0.347	-0.057	0.034	0.035
SQP	0.356	0.087	-0.004	0.148
Eigenvalue	7.600	2.882	1.4534	0.636
Explained variance (%)	58.5	22.2	11.2	4.9
Cumulative variance (%)	58.5	80.7	91.9	96.8