

Additional File 1: Table S1

Supplementary Table S1. Comparison of mean (\pm standard deviation) values of the SF-36 domains according with initial and final phases among patient groups

Variables	Follow-up phases	Group 1^a	Group 2^b
		n = 23	n = 42
SF-36 domains	Initial	Mean \pm SD	p value
	Final	Mean \pm SD	p value
Physical functions	Initial	81 ± 23	0.74
	Final	84 ± 24	
Role-physical	Initial	74 ± 42	0.34
	Final	74 ± 43	
Bodily pain	Initial	70 ± 21	0.19
	Final	64 ± 30	
General health perception	Initial	66 ± 21	0.34
	Final	58 ± 21	
Vitality	Initial	54 ± 26	0.92
	Final	56 ± 24	
Social functioning	Initial	70 ± 32	0.64
	Final	68 ± 31	
Role-emotional	Initial	75 ± 37	0.02
	Final	57 ± 45	
Mental health	Initial	66 ± 23	0.94
	Final	69 ± 22	

^aGroup 1: 23 patients with primary iron overload and homozygosity for the p.Cys282Tyr mutation. ^bGroup 2:

42 patients with primary iron overload and other genotypes: compound heterozygosity for the p.Cys282Tyr/p.His63Asp (n = 10), heterozygosity for the p.Cys282Tyr (n = 4), homozygosity (n = 10) or heterozygosity (n = 8) for the p.His63Asp, or absence of p.Cys282Tyr and p.His63Asp (n = 10).

Additional File 2: Table S2

Supplementary Table S2. Comparison of serum ferritin (ng/mL) values and the SF-36 domains according with initial and final phases in overall patient group

		Serum ferritin initial phase		Serum ferritin finalphase
Physical functions (initial phase,final phase)	r	-0.212	R	-0.352
	p	0.121	P	0.008
Role-physical (initial phase, finalphase)	r	-0.060	R	-0.133
	p	0.661	P	0.332
Bodily pain (initial phase, finalphase)	r	-0.141	R	-0.038
	p	0.305	P	0.784
General health perception(initial phase, final phase)	r	0.008	R	-0.127
	p	0.953	P	0.354
Vitality (initial phase, finalphase)	r	-0.024	R	-0.211
	p	0.864	P	0.121
Social functioning (initial phase,final phase)	r	-0.173	R	-0.200
	p	0.207	P	0.142
Role-emotional (initial phase,final phase)	r	0.157	R	-0.094
	p	0.253	P	0.495
Mental health (initial phase,final phase)	r	0.064	R	0.110
	p	0.643	P	0.424