

Supplementary Material

9p21 Locus Polymorphism is A Strong Predictor of Metabolic Syndrome and Cardiometabolic Risk Phenotypes Regardless of Coronary Heart Disease

Muhammad Mobeen Zafar ¹, Muhammad Saqlain ², Asad Mehmood Raja ¹, Pakeeza Arzoo Shaiq ¹, Muhammad Javaid Asad ¹, Muhammad Kausar Nawaz Shah ³, Farah Fatima ⁴, Hadi Valadi ⁵ Muhammad Nawaz ^{5,*}, Ghazala Kaukab Raja ^{1,*}

¹ University Institute of Biochemistry & Biotechnology, PMAS- Arid Agriculture University Rawalpindi, 46300, Pakistan

² Department of Biochemistry and Molecular Biology, University of Sialkot, Sialkot, 51040, Pakistan

³ Department of Plant Breeding and Genetics, PMAS- Arid Agriculture University Rawalpindi, 46300, Pakistan

⁴ Department of Pathology, Ribeirao Preto Medical School, University of Sao Paulo, Ribeirao Preto, 3900, Brazil

⁵ Department of Rheumatology and Inflammation Research, Institute of Medicine, Sahlgrenska Academy, University of Gothenburg, Gothenburg, 41346, Sweden

* Correspondence: Muhammad Nawaz (muhammad.nawaz@gu.se), Ghazala Kaukab Raja (ghazala@uaar.edu.pk)

Citation: Mobeen Zafar, M.; Saqlain, M.; Mehmood Raja, A.; Arzoo Shaiq, P.; Javaid Asad, M.; Nawaz Shah, M.K.; Fatima, F.; Valadi, H.; Nawaz, M.; Raja, G.K. 9p21 Locus Polymorphism Is A Strong Predictor of Metabolic Syndrome and Cardiometabolic Risk Phenotypes Regardless of Coronary Heart Disease. *Genes* **2022**, *13*, 2226.
<https://doi.org/10.3390/genes1312226>

Academic Editor: Zsolt Ronai

Received: 3 November 2022

Accepted: 23 November 2022

Published: 27 November 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Supplementary table for Figures 3 and 4**Table S1: Comparison of study variables in rs1333049 SNP minor/risk (C) allele and major (G) allele carriers and SD change from population means in C allele carriers**

Study Variables	Non-Carrier (GG)	C Allele Carrier (GC + CC)	SD change from		P-value
			population means in C allele Carrier		
Age (Years)	40.8	46.3	2.3		1.36E-04
Weight (Kg)	70.5	72.3	0.1		3.38E-04
SBP (mm/Hg)	126.2	128.4	-8.4		2.57E-02
DBP (mm/Hg)	82.1	84.6	2.2		1.10E-04
FBG (mg/dl)	95.9	102.4	2.1		3.48E-04
TG (mg/dl)	157	174.3	2.2		3.82E-02
TC (mg/dl)	193.9	215.6	-2.6		2.00E-04
HDL (mg/dl)	48.2	43.9	-1.7		4.55E-03
LDL (mg/dl)	112.1	126.4	1		4.70E-05

SBP; systolic blood pressure, DBP; diastolic blood pressure, FBG; fasting blood glucose, TG; triglycerides, TC; total cholesterol, HDL; high-density lipoprotein, LDL; low density lipoprotein.

Supplementary table for Figures 5 and 6**Table S2: Comparison of adiposity traits in rs1333049 SNP minor/risk (C) allele and major (G) allele carriers and SD change from population means in C allele carriers**

Adiposity Indices	Non-Carrier (GG)	C Allele Carrier (GC + CC)	SD change from		P-value
			population means in C allele Carrier		
BMI (Kg/m ²)	26.2	28.9	0.4		1.10E-04
WC (cm)	86.4	89.4	3.9		3.48E-04
WHR	0.82	0.84	-0.009		4.99E-04
VAI	2.1	2.4	0.1		4.00E-02
BAI	25.3	26.4	1.4		3.30E-02

BMI; body mass index, WC; waist circumference, WHR; waist to hip ratio, VAI; visceral adiposity index, BAI; body adiposity indices,