

Supporting Information

Control of brown rot produced by *Monilinia fructicola* in peaches using a full-spectrum extract of *Zuccagnia punctata* Cav.

Melina G. Di Liberto¹, María Inés Stegmayer², Laura N. Fernández², Ariel D. Quiroga³, Laura A. Svetaz^{1,*} and Marcos G. Derita^{1,2*}

¹ Farmacognosia, Facultad de Ciencias Bioquímicas y Farmacéuticas, Universidad Nacional de Rosario, Suipacha 531, S2002LRK, Rosario, Argentina; mdiliberto@fbioyf.unr.edu.ar

² ICiAgro Litoral, Universidad Nacional del Litoral, CONICET, Kreder 2805, Esperanza, 3080HOF, Argentina; iciagro@santafe-conicet.gov.ar

³ IFISE, Universidad Nacional de Rosario, CONICET, Suipacha 531, S2002LRK, Rosario, Argentina; ifise@rosario-conicet.gov.ar

* Correspondence: lsvetaz@fbioyf.unr.edu.ar; Tel: +54-9341-153-061322 (L.A.S.); mderita@fbioyf.unr.edu.ar, Tel: +54-9341-155-317769 (M.G.D.).

TABLE OF CONTENTS

Figure S1. ^1H NMR (300 MHz) spectrum of 1 in CDCl_3	S3
Figura S2: ^{13}C NMR (300 MHz) spectrum of 1 CDCl_3	S3
Figure S3. EI mass spectrum of 1	S4
Figure S4. ^1H NMR (300 MHz) spectrum of 2 in CDCl_3	S4
Figura S5: ^{13}C NMR (300 MHz) spectrum of 2 CDCl_3	S5
Figure S6. EI mass spectrum of 2	S5




Figura S1: ¹H NMR spectrum of 2',4'-dihydroxy-3'-methoxychalcone (**1**)




Figura S2: ¹³C NMR spectrum of 2',4'-dihydroxy-3'-methoxychalcone (**1**)




Figura S3: EI mass spectrum of 2',4'-dihydroxy-3'-methoxychalcone (**1**)




Figura S4: ^1H NMR spectrum of 2',4'-dihydroxychalcone (**2**)




Figura S5: ^{13}C NMR spectrum of 2',4'-dihydroxychalcone (**2**)




Figura S6: EI mass spectrum of 2',4'-dihydroxychalcone (**2**)