



Influence of the phytosanitary status, cultivar, and harvest time on the phenolic, chlorophyll, and alkaloid content of *Rosa hybrida* L. leaves


Adelina Larisa Munteanu ¹, Loredana Elena Vijan ^{2,*}, Carmen Mihaela Topală ², and Rodi Mitrea ^{1,3}



¹ Doctoral School of Plant and Animal Resources Engineering, University of Craiova, 13 A.I. Cuza Street, RO 200585, Craiova, Romania; adelinamunteanu94@yahoo.com


² Faculty of Sciences, Physical Education and Computer Science, University of Pitesti, 1 Targu din Vale Street, Pitesti, RO 110040, Romania; loredana.vijan@upit.ro



³ Department of Horticulture and Food Science, Faculty of Horticulture, University of Craiova, 13 A.I. Cuza Street, RO 200585, Craiova, Romania; rodi.mitrea@edu.ucv.ro

* Correspondence: loredana.vijan@upit.ro; vloredana2005@yahoo.com







**Melina****Asja**




Figure S1. ATR-FTIR spectra of healthy and affected rose leaves, for 'Orient Express', 'Paul McCartney', 'Terracotta', 'Glorious', 'Mythos', 'Melina', 'Asja', 'Velvet Fragrance', and 'Cluj 2010' cultivars.