

Supplementary file

Novel Sulfamethoxazole Organotin Complexes: Synthesis, Characterization, and Hydrogen Storage Application

Dina S. Ahmed ¹, Noor Emad ², Mohammed Kadhom ³, Emad Yousif ^{2,*} and Mohammed Al-Mashhadani ²

¹ Department of Chemical Industries, Institute of Technology-Baghdad, Middle Technical University, 17564 Baghdad, Iraq; dina_saadi@mtu.edu.iq

² Department of Chemistry, College of Science, Al-Nahrain University, Baghdad 10072, Iraq; noor.emadf@ced.nahrainuniv.edu.iq (N.E.); mohammed.mashhadani@nahrainuniv.edu.iq (M.A.-M.)

³ Department of Environmental Science, College of Energy and Environmental Sciences, Alkarkh University of Science, Baghdad 10081, Iraq; kadhom@kus.edu.iq

* Correspondence: emad.yousif@nahrainuniv.edu.iq




Figure S1. ¹H NMR spectrum of sulfamethoxazole Schiff base (L).




Figure S2. ¹³C NMR spectrum of sulfamethoxazole Schiff base (L).




Figure S3. ¹¹⁹Sn NMR spectrum of Ph₂SnL₂ complex.




Figure S4. ^{119}Sn NMR spectrum of Bu_2SnL_2 complex.




Figure S5. ^{119}Sn NMR spectrum of Me_2SnL_2 complex.