Table S1. Antibiotic susceptibility profiles of lactic acid bacteria isolated from kefir grains

Strain	VAN (30)	ERY (15)	CHL (30)	AMP (10)	PEN (10)	TET (30)	GEN (10)	KAN (30)
Lactobacillus sp. C2K1	22 (S)	15 (I)	39 (S)	29 (S)	24 (S)	0 (R)	11 (S)	9 (R)
Leuconostoc sp. C3K1	0 (R)	30 (S)	46 (S)	32 (S)	39 (S)	0 (R)	0 (R)	0 (R)
Leuconostoc sp. C4K1	0 (R)	31 (S)	46 (S)	38 (S)	30 (S)	0 (R)	0 (R)	0 (R)
Pediococcus sp. C5K1	0 (R)	35 (S)	41 (S)	33 (S)	30 (S)	0 (R)	24 (S)	0 (R)
Leuconostoc sp. C6K1	32 (S)	26 (S)	47 (S)	51 (S)	40 (S)	15 (I)	0 (R)	0 (R)
Leuconostoc sp. C1K2	25 (S)	18 (I)	43 (S)	33 (S)	25 (S)	8 (R)	10 (S)	0 (R)
Lactobacillus sp. C4K2	0 (R)	0 (R)	0 (R)	0 (R)	0 (R)	0 (R)	0 (R)	0 (R)
Lactobacillus sp. C5K2	0 (R)	0 (R)	0 (R)	0 (R)	0 (R)	0 (R)	0 (R)	0 (R)
Lactobacillus sp. C6K2	0 (R)	0 (R)	0 (R)	0 (R)	0 (R)	0 (R)	0 (R)	0 (R)
Lactobacillus sp. C5K3	21 (S)	0 (R)	38 (S)	31 (S)	23 (S)	22 (S)	11 (S)	0 (R)

Note: VAN, vancomycin; ERY, erythromycin; CHL, chloramphenicol; AMP, ampicillin; PEN, penicillin; SXT, trimethoprim–sulfamethoxazole; CLI, clindamycin; OXA, oxacillin; CFZ, cefazolin; CIP, ciprofloxacin; TET, tetracycline; GEN, gentamicin; KAN, kanamycin. R: Resistant; I: Intermediate; S: Susceptible.