**Table S1.** Modified liver staging: architectural changes, fibrosis, and cirrhosis.

| Score | Lobular architecture |
|---|---|
| 0 | Normal (absence of fibrosis) |
| 1 | Fibrous expansion of some portal areas |
| 2 | Fibrous expansion of most portal areas, with portal-portal septa |
| 3 | Fibrous extension of portal spaces with portal-portal and portal-central septa, with possible nodule formation |
| 4 | Cirrhosis with predominant nodular areas in relation to the remaining lobules |

**Table S2.** Modified liver histological activity index (HAI) grading: necro-inflammatory scores.

| **A. Periportal or periseptal interface hepatitis (piecemeal necrosis)** | **Degree** |
|---|---|
| **Absent** | 0 |
| **Mild (focal, few portal areas)** | 1 |
| **Mild/moderate (focal, most portal areas)** | 2 |
| **Moderate (continuous around 60% of tracts or septa)** | 3 |
| **Severe (continuous around >50% of tracts or septa)** | 4 |
| **B. Confluent necrosis** | |
| **Absent** | 0 |
| **Focal confluent necrosis** | 1 |
| **Zone 3 necrosis in some areas** | 2 |
| **Zone 3 necrosis in most areas** | 3 |
| **Zone 3 necrosis+ occasional portal-central (P-C) bridging** | 4 |
| **Zone 3 necrosis+ multiple P-C bridging** | 5 |
| **Panacinar or multiacinar necrosis** | 6 |
| **C. Focal (spotty) lytic necrosis, apoptosis, and focal inflammation** | |
| **Absent** | 0 |
| **One focus or less per 10X objective** | 1 |
| **Two to four foci per 10X objective** | 2 |
| **Five to ten foci per 10X objective** | 3 |
| **More than ten foci per 10X objective** | 4 |
| **D. Portal inflammation** | |
| **Absent of portal lymphocytes** | 0 |
| **Mild number of portal lymphocytes** | 1 |
| **Moderate number of portal lymphocytes** | 2 |
| **Marked number of portal lymphocytes** | 3 |
| **Strongly marked number of portal lymphocytes** | 4 |