

Supplementary figures:

Amyloid Formation in Nanoliter Droplets

Da Yeon Cheong^{1,2,†}, Wonseok Lee^{3,†}, Insu Park⁴, Jinsung Park^{5,*}, and Gyudo Lee^{1,2,*}

¹Department of Biotechnology and Bioinformatics, Korea University, Sejong 30019, Korea

²Interdisciplinary Graduate Program for Artificial Intelligence Smart Convergence Technology, Korea University, Sejong 30019, Korea

³Department of Electrical Engineering, Korea National University of Transportation, Chungju 27469, Korea

⁴Nick Holonyak Jr. Micro and Nanotechnology Laboratory, University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA

⁵Department of Biomechatronic Engineering, College of Biotechnology and Bioengineering, Sungkyunkwan University, Suwon 16419, Korea

*Correspondence: nanojspark@skku.edu (J.P.); lk0807@korea.ac.kr (G.L.)

[†]These authors contributed equally to this work.




Figure S1. Amyloid fibrillation in nanoliter droplets. Bright-field images of nanoliter droplets depend on protein concentration ((a) 3 wt% HEWL (b) 1 wt% HEWL), and corresponding fluorescence images of nanoliter droplets during incubation. The droplet volume for (a) and (b) is 239.7 nL and 237.8 nL, respectively. (Scale bar: 500 μ m).




Figure S2. Droplet shrinking phenomenon. (a) 188.7 nL droplet (b) 147.9 nL droplet.