

Supporting Information
for
Novel azocoumarin derivatives synthesis and characterization

Katarzyna Piechowska¹, Angelika Baranowska-Łączkowska², Krzysztof Z. Łączkowski¹,
 Jolanta Konieczkowska³, Mariola Siwy³, Marharyta Vasylieva³, Paweł Gnida³, Paweł
 Nitschke³, Ewa Schab-Balcerzak³

¹*Department of Chemical Technology and Pharmaceuticals, Faculty of Pharmacy, Collegium Medicum, Nicolaus Copernicus University, Jurasza 2, 85-089 Bydgoszcz, Poland; kpiechowska@cm.umk.pl (K.P); krzysztof.laczkowski@cm.umk.pl (K.Z.L.)*

²*Institute of Physics, Kazimierz Wielki University, Powstańców Wielkopolskich 2, 85-090, Bydgoszcz, Poland; angelika.baranowska@ukw.edu.pl (A.B.L.)*

³*Centre of Polymer and Carbon Materials, Polish Academy of Sciences, 34 M. Curie-Skłodowska Str., 41-819 Zabrze, Poland*

Table S1. Solubility of azocoumarin derivatives in different solvents (2 mg of the compound in 1ml of solvent).

Sample	H ₂ O		THF		CHCl ₃		DMF		CH ₃ OH	
	room temp.	boiling temp.	room temp.	boiling temp.	room temp.	boiling temp.	room temp.	boiling temp.	room temp.	boiling temp.
6-a	–	+/-	+/-	+/-	+/-	+/-	+	+	+/-	+/-
6-b	–	–	+/-	+/-	+/-	+/-	+	+	+/-	+/-
6-c	–	+/-	+/-	+/-	+/-	+/-	+	+	+/-	+/-
6-d	–	+/-	+/-	+/-	+/-	+/-	+	+	+/-	+/-
6-e	–	–	+/-	+/-	+/-	+/-	+	+	+/-	+/-
6-f	–	–	+/-	+/-	+/-	+/-	+/-	+	+/-	+/-
6-g	+/-	+/-	+/-	+/-	+/-	+/-	+/-	+	+/-	+/-
6-h	–	+/-	+/-	+/-	+	+	+/-	+	+/-	+/-
6-i	–	+/-	+/-	+/-	+	+	+/-	+	+/-	+/-




Figure S1. UV-vis spectra of **6b** registered before and after irradiation ($\lambda=365\text{nm}$, 2.9 W , 30s) in (a) chloroform, (b) ethanol, (c) DMF, and (d) polystyrene.






Figure S2. Absorbance spectra of compound 6a-6i in DCM

Table S2. Content of *cis*-isomer directly after the 5 min of UV-light irradiation, and time of *cis* to *trans* transformation of dyes in polystyrene matrix.



Code	Isosbestic points [nm]	Content of <i>cis</i> isomer (%)	Time of <i>cis-trans</i> conversion [days]
6a+PS	418	13	8
6b+PS	420	14	8
6c+PS	421	13	>8
6d+PS	424	11	>8
6f+PS	419	14	7
6g+PS	410	14	>8
6h+PS	437	7	7
6i+PS	407	22	>8

¹H, ¹³C NMR and HRMS spectra of compounds **3-5**, **6a-6i** and additional DEPT-135, COSY, NOESY, HMBC, HSQC spectra for compound **6g**.



Compound 3




Compound 5






Compound 6a








Compound 6b







Compound 6c






Compound 6d






210322_6dA 33 (0.357) Cm (33:40-(4:8+82:97))

TOF MS ES+
2.35e5




Compound 6e





210322_6i_A 17 (0.197) Cm (17:20-(3:8+81:97))

TOF MS ES+
5.16e6




Compound 6f




210322_6fA 24 (0.256) Cm (24:41-(4:8+87:97))


TOF MS ES+
1.19e6




Compound 6g



COSY




HMBC




HSQC

6g
K.Laczkowski




NOESY

6g
K.Laczkowski





210322_6eA 33 (0.357) Cm (33:40-3:8)

TOF MS ES+
1.45e6





Compound 6h





Compound 6i





210322_6hA 29 (0.311) Cm (29:40-3:8)

