



Supplemental table S5

Temperature-biased miRNA expression patterns during European sea bass (*Dicentrarchus labrax*) development.

Maria Papadaki^{1,2}, Elisavet Kaitetzidou¹, Ioannis Papadakis¹, Dimitris Sfakianakis², Papandroulakis N.¹, Constantinos C. Mylonas¹, Elena Sarropoulou^{1*}

| | Stage | Mouth opening | First feeding | Flexion | All fins | Metamorphosis |
|---------|------------------|---------------|---------------|--------------|--------------|---------------|
| 20 °C | Age (dph) | 3 | 6 | 18 | 40 | 53 |
| | Mean length (mm) | 5,17 ± 0.14 | 5,53 ± 0.12 | 10,47 ± 1.19 | 21 ± 0.5 | 29,67 ± 3.43 |
| 17,5 °C | Age (dph) | 5 | 9 | 20 | 47 | 62 |
| | Mean length (mm) | 5,53 ± 0.08 | 5,86 ± 0.15 | 10,94 ± 0.7 | 20,63 ± 0.79 | 27,35 ± 1.46 |
| 15°C | Age (dph) | 6 | 11 | 30 | 53 | 86 |
| | Mean length (mm) | 5,41 ± 0.11 | 5,78 ± 0.13 | 12,09 ± 0.96 | 19,25 ± 0.76 | 29,10 ± 2.97 |