

Figure S1. Distribution of CFU per gram of tissue values obtained for strain CA1. Control is referred to as a positive control (untreated biofilm of *C. albicans*). Biofilms were treated with: AmfB – amphotericin B (50 μg/mL), lipopeptide L1 at a concentration equal to MBEC, ½ MBEC and FIC values (MBEC – minimum biofilm eradication concentration; FIC – fractional inhibitory concentration). NS – non significant results ($p>0.05$), */** - results with statistical significance ($p<0.05$).



Figure S2. Distribution of CFU per gram of tissue values obtained for strain CA1. Control is referred to as a positive control (untreated biofilm of *C. albicans*). Biofilms were treated with: AmfB – amphotericin B (50 μg/mL), lipopeptide C1 at a concentration equal to MBEC, ½ MBEC and FIC values (MBEC – minimum biofilm eradication concentration; FIC – fractional inhibitory concentration). NS – non significant results ($p>0.05$), */** - results with statistical significance ($p<0.05$).



Figure S3. Distribution of CFU per gram of tissue values obtained for strain CA2. Control is referred to as a positive control (untreated biofilm of *C. albicans*). Biofilms were treated with: AmfB – amphotericin B (50 μg/mL), lipopeptide L1 at a concentration equal to MBEC, ½ MBEC and FIC values (MBEC – minimum biofilm eradication concentration; FIC – fractional inhibitory concentration). NS – non significant results ($p>0.05$), */** - results with statistical significance ($p<0.05$).



Figure S4. Distribution of CFU per gram of tissue values obtained for strain CA2. Control is referred to as a positive control (untreated biofilm of *C. albicans*). Biofilms were treated with: AmfB – amphotericin B (50 μg/mL), lipopeptide C1 at a concentration equal to MBEC, ½ MBEC and FIC values (MBEC – minimum biofilm eradication concentration; FIC – fractional inhibitory concentration). NS – non significant results ($p>0.05$), */** - results with statistical significance ($p<0.05$).