



Supplementary

Phosphorylation of CaMK and CREB-Mediated Cardiac Aldosterone Synthesis Induced by Arginine Vasopressin in Rats with Myocardial Infarction

Yuan-Sheng Zhai ^{1,2,†}, Jie Li ^{1,2,†}, Longyun Peng ^{1,2}, Guihua Lu ^{1,2}, and Xiuren Gao ^{1,2,*}

¹ Department of Cardiology, the First Affiliated Hospital, Sun Yat-Sen University, Guangzhou 510800, China

² Key Laboratory on Assisted Circulation, Ministry of Health, Guangzhou 510800, China

† These authors contribute equally to the work.

* Correspondence: gaoxr@mail.sysu.edu.cn; Tel.: +86-020-82394178




Figure S1. Representative M-mode echocardiographic images of rats. 16w sham, 16-week post-sham-operation group; 6w MI, 6-week post-infarction group; 11w MI, 11-week post-infarction group; 16w MI, 16-week post-infarction group.




Figure S2. Correlation analyses among the rats. LVEF, left ventricular ejection fraction. A: Correlation analysis between aldosterone and type I collagen expression. B: Correlation analysis between aldosterone and LVEF. C: Correlation analysis between AVP and type I collagen expression. D: Correlation analysis between AVP and LVEF.