

Inhibition of soluble epoxide hydrolase activity by components of *Glycyrrhiza uralensis*

Jang Hoon Kim ¹, Yun-Chan Huh ¹, Mok Hur ¹, Woo Tae Park ¹, Youn-Ho Moon ¹, Tae Il Kim ¹, Yong Il Kim ¹, Seon Mi Kim ¹, Jeonghoon Lee ^{1,*} and Ik Soo Lee ^{2,*}

¹ Department of Herbal Crop Research, National Institute of Horticultural and Herbal Science, RDA, Eumseong 27709, Republic of Korea

² Km Convergence Research Division, Korea Institute of Oriental Medicine, Daejeon 34134, Republic of Korea

* Correspondence: artemisia@korea.kr (J.L.); knifer48@kiom.re.kr (I.S.L.);
Tel.: +82-42-871-5670 (J.L.); +82-42-868-9462 (I.S.L.)




Figure S1. ¹H NMR(A) and ¹³C NMR(B) spectra of compound 1 in MeOD-*d*₄ (400 MHz for ¹H NMR, 100 MHz for ¹³C NMR)




Figure S2. LC-MS(A), ^1H NMR(B) and ^{13}C NMR(C) spectra of compound **2** in $\text{MeOD-}d_4$ (400 MHz for ^1H NMR, 100 MHz for ^{13}C NMR)



Figure S3. ^1H NMR(A) and ^{13}C NMR(B) spectra of compound **3** in $\text{MeOD-}d_4$ (400 MHz for ^1H NMR, 100 MHz for ^{13}C NMR)




Figure S4. LC-MS(A), ^1H NMR(B) and ^{13}C NMR(C) spectra of compound **4** in $\text{MeOD-}d_4$ (400 MHz for ^1H NMR, 100 MHz for ^{13}C NMR)




Figure S5. LC-MS(A), ^1H NMR(B) and ^{13}C NMR(C) spectra of compound **5** in $\text{MeOD-}d_4$ (400 MHz for ^1H NMR, 100 MHz for ^{13}C NMR)




Figure S6. LC-MS(A), ^1H NMR(B) and ^{13}C NMR(C) spectra of compound **6** in $\text{MeOD-}d_4$ (400 MHz for ^1H NMR, 100 MHz for ^{13}C NMR)




Figure S7. LC-MS(A), ^1H NMR(B) and ^{13}C NMR(C) spectra of compound **7** in $\text{MeOD-}d_4$ (400 MHz for ^1H NMR, 100 MHz for ^{13}C NMR)




Figure S8. LC-MS(A), ^1H NMR(B) and ^{13}C NMR(C) spectra of compound **8** in $\text{MeOD-}d_4$ (400 MHz for ^1H NMR, 100 MHz for ^{13}C NMR)




Figure S9. ^1H NMR(A) and ^{13}C NMR(B) spectra of compound **9** in $\text{MeOD-}d_4$ (400 MHz for ^1H NMR, 100 MHz for ^{13}C NMR)




Figure S10. LC-MS(A), ^1H NMR(B) and ^{13}C NMR(C) spectra of compound **10** in $\text{MeOD-}d_4$ (400 MHz for ^1H NMR, 100 MHz for ^{13}C NMR)




Figure S11. LC-MS(A), ¹H NMR(B) and ¹³C NMR(C) spectra of compound **11** in MeOD-*d*₄ (400 MHz for ¹H NMR, 100 MHz for ¹³C NMR)