

## Supporting Information


# Multifunctional oxidized dextran as matrix for stabilization of octahedral molybdenum and tungsten iodide clusters in aqueous media

Ekaterina V. Pronina <sup>1</sup>, Yuri A. Vorotnikov <sup>1,\*</sup>, Tatiana N. Pozmogova <sup>1</sup>, Alphiya R. Tsygankova <sup>1</sup>, Kaplan Kirakci <sup>2</sup>, Kamil Lang <sup>2</sup>, Michael A. Shestopalov <sup>1</sup>

**Table S1.** Chemical groups content and molecular weights of initial and oxidized dextrans [1].

Sample	Carbonyl groups, mmol·g <sup>-1</sup>	Acidic groups, mmol·g <sup>-1</sup>	M <sub>w</sub> , kDa
Dex6	-	-	6.2
NaOxDex6(1)	0.6	1.4	3.8
NaOxDex6(2)	0.6	2.3	3.9
Dex60	-	-	60.2
NaOxDex60(1)	1.0	1.0	14.0
NaOxDex60(2)	1.3	1.5	4.5
NaOxDex60(2)-red	0.8	1.5	4.5


1. Pronina, E.V.; Vorotnikov, Y.A.; Pozmogova, T.N.; Solovieva, A.O.; Miroshnichenko, S.M.; Plyusnin, P.E.; Pishchur, D.P.; Eltsov, I.V.; Edeleva, M.V.; Shestopalov, M.A.; et al. No Catalyst Added Hydrogen Peroxide Oxidation of Dextran: An Environmentally Friendly Route to Multifunctional Polymers. *ACS Sustain. Chem. Eng.* **2020**, *8*, 5371-5379, doi:10.1021/acssuschemeng.0c01030.




**Figure S1.** FTIR spectra of OxDexQ(n), NaOxDexQ(n), Mo<sup>100</sup>@NaOxDexQ(n), and W<sup>100</sup>@NaOxDexQ(n) (A – Q = 6, n = 1; B – Q = 6, n = 2; C – Q = 60, n = 2).

**Table S2.** {M<sub>6</sub>I<sub>8</sub>} content in materials (μmol·g<sup>-1</sup>) according to ICP-AES

x	10	50	100
<b>M = Mo</b>			
Dex6	-	-	28
NaOxDex6(1)	33	122	139
NaOxDex6(2)	31	127	144
Dex60	-	-	54
NaOxDex60(1)	31	97	115
NaOxDex60(2)	30	118	130
<b>M = W</b>			
Dex6	-	-	48
NaOxDex6(1)	11	51	82
NaOxDex6(2)	13	51	87
Dex60	-	-	59
NaOxDex60(1)	20	49	125
NaOxDex60(2)	16	58	119




**Figure S2.** UV-vis spectra of Mo<sup>100</sup>@Dex60 (A), Mo<sup>100</sup>@NaOxDex60(1) (B), and Mo<sup>100</sup>@NaOxDex60(2) in water over time.




**Figure S3.** UV-vis spectra of M<sup>100</sup>@NaOxDex60(1), M = Mo (A) and W (B) in water, in comparison with corresponding initial clusters.

**Table S3.** Formulation of culture medium DMEM.


Component	Concentration, mg/L	Component	Concentration, mg/L
<b>Inorganic salts</b>		<b>Amino acids</b>	
Calcium Chloride · 2H <sub>2</sub> O	264.92	L-Arginine · HCl	84.00
Ferric Nitrate · 9H <sub>2</sub> O	0.10	L-Cystine · 2HCl	62.58
Magnesium Sulfate (Anhydr.)	97.67	L-Glutamine	584.00
Potassium Chloride	400.00	Glycine	30.00
Sodium Chloride	6400.00	L-Histidine · HCl · H <sub>2</sub> O	42.00
Sodium Phosphate, Monobasic (Anhydr.)	108.69	L-Isoleucine	104.80
<b>Vitamins</b>		L-Leucine	104.80
Choline Chloride	4.00	L-Lysine · HCl	146.20
Folic Acid	4.00	L-Methionine	30.00
myo-Inositol	7.00	L-Phenylalanine	66.00
Nicotinamide	4.00	L-Serine	42.00
D-Pantothenic Acid, Hemicalcium Salt	4.00	L-Threonine	95.20
Pyridoxal · HCl	4.00	L-Tryptophan	16.00
Riboflavin	0.40	L-Tyrosine · Na <sub>2</sub> · 2H <sub>2</sub> O	103.79
Thiamine · HCl	4.00	L-Valine	93.60
<b>Other components</b>			
D-Glucose	1000.00	Sodium Bicarbonate	3700.00



**Figure S4.** UV-vis spectra of Mo<sup>x</sup>@NaOxDex60(*n*) (A – *x* = 10, *n* = 1; B – *x* = 10, *n* = 2; C – *x* = 50, *n* = 1; D – *x* = 50, *n* = 2) in DMEM culture medium over time.



**Figure S5.** UV-vis spectra of  $W^{100}@NaOxDexQ(n)$  (A – Q = 6, n = 1; B – Q = 6, n = 2; C – Q = 60, n = 1; D – Q = 60, n = 2) in DMEM culture medium over time.



**Figure S6.** UV-vis spectra of  $W^x@NaOxDex60(n)$  (A –  $x = 10$ ,  $n = 1$ ; B –  $x = 10$ ,  $n = 2$ ; C –  $x = 50$ ,  $n = 1$ ; D –  $x = 50$ ,  $n = 2$ ) in DMEM culture medium over time.




Figure S7. FTIR spectra of NaOxDex60(2) and NaOxDex60(2)-red.




Figure S8. Emission spectra of fresh and 8-days old solution of  $\text{Mo}^{50}\text{@NaOxDex60(1)}$  in PBS (air and Ar) (A); Emission spectra of fresh and 8-days old solution of  $\text{W}^{50}\text{@NaOxDex60(1)}$  in PBS (air and Ar) (B).