

Supplementary Material

Affinity resins for isolation of immunoglobulins G obtained via biocatalytic technology

Mikhail N. Tereshin ^{1,2}, Tatjana D. Melikhova ¹, Barbara Z. Eletskaya ¹, Olga B. Ksenofontova ¹, Pavel V. Pantyushenko ¹, Maria Ya. Berzina ^{1,*}, Igor Ivanov ², Igor V. Myagkikh ¹ and Vasilijs N. Stepanenko ^{1,2}


¹ M.M. Shemyakin and Yu.A. Ovchinnikov Institute of Bioorganic Chemistry, Russian Academy of Sciences (IBCh RAS), Miklukho-Maklaya street, 16/10, 117437, Moscow, Russia; misha060596@gmail.com (M. N. T.); tdm-63@yandex.ru (T. D. M); fraubarusya@gmail.com (B. Z. E.); labpeptos@gmail.com (O. B. K); pantyushenko.pavel@gmail.com (P. V. P.); berzina_maria@mail.ru (M. Ya. B.); myagkikh@ibch.ru (I. V. M.); svn@ibch.ru (V. N. S.)

² Lomonosov Institute of Fine Chemical Technologies, MIREA - Russian Technological University, Vernadskogo pr. 86, 119571 Moscow, Russia; igor_ivanov@gmx.de

Table of Content

¹ H spectrum of compound Boc-Gly-Gly-Gly-Gly-OH (9)	S1
¹ H spectrum of compound Boc-Gly-Gly-EDA-Gly-Gly-Boc (12)	S2

SI-1 Boc-Gly₄-OH



Chemical Formula: C₁₃H₂₂N₄O₇

Molecular Weight: 346,34





Figure S1. The ¹H NMR spectrum of Boc-Gly₄-OH

SI-2 Boc-Gly-Gly-EDA-Gly-Gly-Boc



Chemical Formula: C₂₀H₃₆N₆O₈

Exact Mass: 488,26




Figure S2. The ¹H NMR spectrum of Boc-Gly-Gly-EDA-Gly-Gly-Boc