



**Figure S1.** Frequency (%) of symptoms reported by Bichara et al. [8]. LR (Long Response) frequencies were obtained from the Bichara´s paper Table 5 (Persistent) and the SR (Short Response) were obtained from the sum of Table 5's Nonpersistent plus the Nonreactive from Bichara´s paper Table 2. (Wilcoxon paired test; Z=3.23; p=0.0012).

**Table S1.** COVID-19 Symptoms in Belém, Pará, Brazil population.

| **Severity** | **Nonsevere** | **Severe** | **Non-Hospitalized** | **Hospitalized** |
|---|---|---|---|---|
| **Sample Sizes** | N | N | N | N |
| **References** | Sarges et al.[18] | | Rodrigues et al.[19] | |
| **Symptoms** | | | | |
| **Fever** | 69.33 | 73.61 | 61.05 | 75 |
| **Cough** | 62 | 84.72 | 48.42 | 81.81 |
| **Runny nose/Coryza** | 40.67 | 33.33 | 40 | 35.61 |
| **Headache** | 58 | 58.33 | 61.05 | 53.79 |
| **Sore throat** | 40 | 31.94 | 42.1 | 32.57 |
| **Chest pain** | 40 | 52.78 | 31.58 | 42.42 |
| **Abdominal pain** | 16.67 | 30.56 | 16.84 | 23.48 |
| **Muscle or body pain** | 58.67 | 58.33 | 53.68 | 59.85 |
| **Nausea** | 19.33 | 37.5 | 17.89 | 34.09 |
| **Vomiting** | 10 | 15.28 | | |
| **Diarrhea** | 42.67 | 44.44 | 34.74 | 47.73 |
| **Dyspnea** | 36.67 | 73.61 | 29.47 | 60.61 |
| **Weakness** | 49.33 | 65.28 | 20 | 27.27 |
| **Fatigue** | 54.67 | 75 | | |
| **Anosmia** | 52.67 | 38.89 | 63.16 | 39.39 |
| **Ageusia** | 50.67 | 38.89 | | |