

Figure S1. The mRNA expression levels of clusterin, CysC, GSTπ and TIMP-1 were measured in RPTEC/TERT1 cell treated with TGF-β1 with/without i-bodies. Results are presented as mean±SEM. n = 4.

Table S1. The sequence of primers used for quantitative RT-PCR.

Species	Target	Forward (5'-3')	Reverse (5'-3')
Human	**GAPDH**	AGCCACATCGCTCAGACAC	GCCCAATACGACCAAATCC
	TIMP-1	GGGGCTTCACCAAGACCTAC	GGAAGCCCTTTTCAGAGCCT
	CysC	GCCTGTGCCTATCACCTCTTAT	CCTTCTCTGTCTGTCTCCTGGT
	Clusterin	CCAGGACAGGTTCTTCACCC	CGTACGGAGAGAAGGGCATC
	GST-π	TATTTCCCAGTTCGAGGCCG	TACAGGGTGAGGTCTCCGTC
Mouse	**β-Actin**	CAGCTGAGAGGGAAATCGTG	CGTTGCCAATAGTGATGACC
	COL-3	TCCCCTGGAATCTGTGAATC	TGAGTCGAATTGGGGAGAAT
	α-SMA	CATCTTTCATTGGGATGGAG	TTAGCATAGAGATCCTTCCTG