

Population Density and Host Preference of the Japanese Pine Sawyer (*Monochamus alternatus*) in the Qinling–Daba Mountains of China

Junke Nan ^{1,2}, Jingyu Qi ², Yuexiang Yang ², Mengqin Zhao ², Chaoqiong Liang ³, Hong He ^{2,*} and Cong Wei ^{1,*}

¹ Key Laboratory of Plant Protection Resources and Pest Management of the Ministry of Education, College of Plant Protection, Northwest A&F University, Yangling 712100, China

² Key Laboratory of National Forestry and Grassland Administration for Control of Forest Biological Disasters in Western China, College of Forestry, Northwest A&F University, Yangling 712100, China

³ Shaanxi Academy of Forestry, Xi'an 710082, China

Supplementary figure:




Figure S1. Frequency distribution of head capsule width and pronotum width of *Monochamus alternatus* larvae. A: Frequency distribution of head capsule width of *M. alternatus* larvae on *P. tabuliformis*, *P. armandii* and *P. massoniana*; B: Frequency distribution of pronotum width of *M. alternatus* larvae on *P. tabuliformis*, *P. armandii* and *P. massoniana*.

Supplementary table captions:

Table S1. Different groups were divided according to the head capsule width of *Monochamus alternatus* larvae.

Table S2. Different groups were divided according to the pronotum width of *Monochamus alternatus* larvae.

Table S3. Feeding and oviposition preference of *Monochamus alternatus* on three pine trees.