

**Table S1.** Generalized Linear Model analyses of the number consumed whiteflies and aphids, comparing among whitefly densities offered, inside each treatment of alternative prey (aphids) for Experiment 1 (without intraguild predation) and 2 (with intraguild predation using parasitized whitefly nymphs).

Consumed whiteflies among offered whitefly densities							
Experiment	Treatment	GLM family	Df	Deviance	Resid. Df	Resid. Dev	P
Without IGP	No AP	Negative Binomial	4	85.57	25	33.696	< 0.0001
	5_AP	Negative Binomial	4	30.477	25	35.37	< 0.0001
	25_AP	Negative Binomial	4	34.651	25	35.316	< 0.0001
	80_AP	Negative Binomial	4	26.165	25	32.094	< 0.0001
With IGP	No AP	Negative Binomial	4	108.12	25	28.359	< 0.0001
	5_AP	Negative Binomial	4	69.216	25	28.752	< 0.0001
	25_AP	Negative Binomial	4	79.94	25	31.722	< 0.0001
	80_AP	Negative Binomial	4	25.259	25	30.588	< 0.0001
Consumed aphids among offered whitefly densities							
Without IGP	5_AP	Gaussian	4	0.33333	25	16.333	0.9725
	25_AP	Poisson	4	1.8037	25	73.142	0.7718
	80_AP	Gaussian	4	912.13	25	4610.2	0.2929
With IGP	5_AP	Gaussian	4	8.5333	25	35.333	0.1963
	25_AP	Gaussian	4	371.33	25	416.67	0.0002
	80_AP	Negative Binomial	4	4.2927	25	30.497	0.3678