Article

Light Intensity Modulates Locomotor Behavior and Predation in Different Color Morphs of the Harlequin Ladybird, *Harmonia axyridis*

Xing-Xing Wang [1], Ya-Nan Liu [1], Chun-Yan Huang [1], Rui-Yan Li [1], Zhi-Wei Jiang [1], Chen-Yang Liu [1], Tong-Xian Liu [2] and Yi Zhang [1,*]

[1] Shandong Engineering Research Center for Environmentally Friendly Agricultural Pest Management, College of Plant Health and Medicine, Qingdao Agricultural University, Qingdao 266000, China; wxx0521@qau.edu.cn (X.-X.W.); lyn2001@stu.qau.edu.cn (Y.-N.L.); 20220205470@stu.qau.edu.cn (C.-Y.H.); lry027@stu.qau.edu.cn (R.-Y.L.); 15562731283@163.com (Z.-W.J.); 13613128117@163.com (C.-Y.L.)

[2] Institute of Entomology, Guizhou University, Guiyang 550025, China; tx.liu@gzu.edu.cn

* Correspondence: yizhang085@qau.edu.cn

Table S1. List of primers utilized in this study.

	Forward (5' to 3')	Reverse (5' to 3')	PCR
Q-*UVSop-1*	GCCAGCTGTATAGATCCCTT	AGTTTGAATTGGCATCATCG	RT-qPCR
Q-*UVSop-2*	TCAAAGAAGTTTGGACATGC	CCATCCGAGTCTATGAAGAA	RT-qPCR
Q-*HaLWSop*	TAGGTATAGGAATTCTGGGCTT	AAATGGATTCCCAAAGATGGA	RT-qPCR
Q-*HaRps3* (Ref)	GGCTACCAGAACCGAACAGAG	GTGCTATGGCGCATAATCCT	RT-qPCR
ds-*UVSop-1*	T7+GGCTCCCAAAGAGAGAAACGA	T7+GCCAACCTCCTTATGCTGGA	dsRNA
ds-*UVSop-2*	T7+AATGGGTTCGGGGGAAAATGA	T7+GTTGGAAGCTGTTCTCAGGC	dsRNA
ds-*HaLWSop*	T7+GTGTGCAGTGACTTTTCCGT	T7+GGGGGTACTGTTAGAGCCAA	dsRNA
ds-*Muslta*	T7+CACCCTCTCCACGAATTG	T7+TAGAAGATGCTGCTGTTTCA	dsRNA