Table S1. Nymphal mortality of *Myzus persicae* after a single application of the essential oil to newly emerged first-instar nymphs (N1) at different concentrations (μL mL^{-1}).

Concentration (μL mL^{-1})	L1	L2	L3	L4	Cumulative mortality (N1 → adult)	Adult emergence
Control	0/30 (0.0)	0/30 (0.0)	0/30 (0.0)	0/30 (0.0)	0/30 (0.0%)	30/30 (100%)
0.5	0/30 (0.0)	0/30 (0.0)	0/30 (0.0)	0/30 (0.0)	0/30 (0.0%)	30/30 (100%)
1	0/30 (0.0)	0/30 (0.0)	0/30 (0.0)	0/30 (0.0)	0/30 (0.0%)	30/30 (100%)
2	0/30 (0.0)	0/30 (0.0)	0/30 (0.0)	2/30 (6.7)	2/30 (6.7%)	28/30 (93.3%)
4	0/30 (0.0)	0/30 (0.0)	0/30 (0.0)	3/30 (10.0)	3/30 (10.0%)	27/30 (90.0%)
8	0/30 (0.0)	2/30 (6.7)	2/28 (7.1)	1/26 (3.8)	5/30 (16.7%)	25/30 (83.3%)
12	3/30 (10.0)	5/27 (18.5)	3/22 (13.6)	4/19 (21.1)	15/30 (50.0%)	15/30 (50.0%)

Values for each instar are the number of nymphs that died at that instar / the number of nymphs that entered it (mortality, %). Cumulative mortality and adult emergence are relative to the 30 N1 initially treated per concentration. Nymphs were treated once (N1) and then monitored individually on fresh, untreated leaves.