ClinicalTrials.gov Search Results 02/01/2025

	Title	Status	Study Results	Conditions	Interventions	Locations
1	Comparison of LAA-Closure vs Oral Anticoagulation in Patients With NVAF and Status Post Intracranial Bleeding.	Recruiting	No Results Available	• Atrial Fibrillation (AF) • Intracranial Bleed	• Device: Percutaneous closure of the LAA (Watchman / Watchman FLX)	• Universitätsherzzentrum Freiburg - Bad Krozingen, Freiburg, Baden Württemberg, Germany • Klinikum Friedrichshafen GmbH, Friedrichshafen, Baden-Wurttemberg, Germany • Universitätsklinikum der J.W. Goethe-Universität Frankfurt, Frankfurt am main, Hessen, Germany • Knappschaftskrankenhaus Bottrop Gmbh, Bottrop, Nordrhein Westfahlen, Germany • Klinikum Chemnitz, Chemnitz, Sachsen, Germany • Städtisches Klinikum Friedrichstadt Dresden, Dresden, Sachsen, Germany • Klinikum St. Georg gGmbH, Leipzig, Sachsen, Germany • HBK Zwickau, Zwickau, Sachsen, Germany • Katholisches Krankenhaus "St. Johann Nepomuk", Erfurt, Thüringen, Germany • SRH Wald-Klinikum Gera GmbH, Gera, Thüringen, Germany • and 32 more
2	Prevention of Stroke by Left Atrial Appendage Closure in Atrial Fibrillation Patients After Intracerebral Hemorrhage	Recruiting	No Results Available	• Atrial Fibrillation • Stroke • Intracerebral Hemorrhage	• Device: LAAO • Drug: Medical Therapy	• Aarhus University Hospital, Aarhus, Denmark • Bispebjerg University Hospital, Bispebjerg, Denmark • Rikshospitalet Glostrup, Glostrup, Denmark • Herlev sjukhus, Herlev, Denmark • Odenses Universitetssjukhus, Odense, Denmark • Helsinki University Hospital, Helsinki, Finland • North Karelia Central Hospital, Joensuu, Finland • Kuopio University Hospital, Kuopio, Finland • Turku University Hospital, Turku, Finland • Vaasa Centralsjukhus, Vaasa, Finland • and 7 more

U.S. National Library of Medicine | U.S. National Institutes of Health | U.S. Department of Health & Human Services