

Table S1. Prevalence of uncommon clinical features in the present series of KBG syndrome patients.

Clinical features	P1	P2	P3	P4	P5	P6	P7	P8	P9	P10	P11	P12	P13	P14	P15	P16	P17	P18	P19	P20	P21	P22	P23	P24
Sex	M	F	M	M	F	M	M	F	F	M	F	F	M	M	M	F	M	F	F	M	M	M	F	M
Neuropsychological anomalies																								
Epilepsy	-	-	-	+	-	+	-	-	-	+	+	-	-	-	+	-	-	+	-	+	-	-	-	-
Motor tics	-	-	-	-	-	-	-	-	-	-	-	-	-	+	-	-	-	-	-	-	-	-	-	-
Cerebral malformation	-	n.e.	n.e.	-	+	-	-	n.e.	n.e.	-	-	n.e.	+	-	n.e.	-	n.e.	n.e.	n.e.	n.e.	-	-	+	-
Ocular anomalies																								
Strabismus	+	-	-	-	-	+	-	-	-	-	-	-	+	+	-	+	-	-	-	-	-	-	-	-
Otorhinolaringological anomalies																								
Hearing loss	-	-	-	-	-	-	-	-	-	-	-	-	+	-	-	-	-	-	+	-	-	+	-	-
Sensorineural	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Conductive	-	-	-	-	-	-	-	-	-	-	-	-	+	-	-	-	-	-	+	-	-	+	-	-
Gastroenterological problems																								
Gastroesophageal reflux	+	-	-	+	-	-	-	-	-	-	+	-	+	+	-	+	-	-	-	-	-	-	+	-
Treatments medication at the time of assessment																								
	-	-	-	+	-	+	-	-	-	+	+	-	-	-	+	-	-	-	-	-	-	-	-	-
				levetiracetam		risperdal venlafaxina valproic acid				levetiracetam	carbamazepine				valproic acid									

P, patient; F, female, M, male; n.e., not evaluable, + positive, - negative, *P6 takes drugs because he also had a behavioural disorder