

## *Supplementary Material*

## 1 Supplementary Tables

### **1.1 Supplementary Table S1. Represented z-value among significant cluster between verum and sham acupuncture intervention**

	Verum (N=15)			Sham(N=12)		
	Pre	During	Post	Pre	During	Post
<b>Post vs Pre</b>						
R Mid Frontal G	-0.07±0.12	-0.10±0.11	-0.09±0.15	-0.05±0.12	-0.07±0.11	-0.14±0.16
<b>During vs Pre</b>						
R Inf Parietal Lob	-0.04±0.10	-0.05±0.11	-0.03±0.15	-0.01±0.10	-0.04±0.15	-0.06±0.14
R Mid Occipital G	-0.06±0.09	-0.14±0.12	-0.03±0.11	-0.04±0.10	-0.03±0.09	-0.05±0.14
<b>Post vs During</b>						
L dACC	0.03±0.15	0.10±0.15	0.04±0.18	-0.02±0.10	0.11±0.13	-0.02±0.13
L Postcentral G	-0.01±0.14	0.01±0.10	0.01±0.18	0.05±0.23	-0.08±0.29	0.03±0.12

Pre: pre-treatment; During: during-treatment; Post: post-treatment; L: left; R: right; Mid: middle; dACC: dorsal anterior cingulate cortex; G: gyrus; Lob: lobule. All values are presented as mean  $\pm$  SD.

## **1.2 Supplementary Table S2. Changes in functional connectivity map after verum or sham acupuncture intervention**

R Cuneus	19/18	503	3.55	2	-78	32	n.s.
L Precuneus	19		3.24	-8	-82	44	
<b>Post &lt; During</b>							
L Postcentral G	43	595	4.21	-64	-8	20	R Mid Frontal G
L Sup Temporal G	42		3.87	-66	-26	14	R Sup Frontal G
L Mid Frontal G	9		3.81	-54	18	36	

BA: Brodmann area; Size: number of voxels in the cluster; t-max: peak t value; Pre: pre-treatment; During: during-treatment; Post: post-treatment; L: left; R: right; Bil: bilateral; Sup: superior; Mid: middle; Inf: inferior; ACC: anterior cingulate cortex; G: gyrus; Lob: lobule; n.s.: non-significant.