

Cardiopulmonary Exercise Performance and Endothelial Function in Convalescent COVID-19 Patients


Pasquale Ambrosino ^{1,*,†}, **Paolo Parrella** ^{2,‡}, **Roberto Formisano** ¹, **Giovanni Perrotta** ¹, **Silvestro Ennio D'Anna** ³, **Marco Mosella** ³, **Antimo Papa** ^{1,‡} and **Mauro Maniscalco** ^{3,*‡}

- ¹ Istituti Clinici Scientifici Maugeri IRCCS, Cardiac Rehabilitation Unit of Telese Terme Institute, 82037 Telese Terme, Italy; roberto.formisano@icsmaugeri.it (R.F.); giovanni.perrotta@icsmaugeri.it (G.P.); antimo.papa@icsmaugeri.it (A.P.)
- ² Ospedale Sacro Cuore di Gesù Fatebenefratelli, 82100 Benevento, Italy; paoloparrella01@libero.it
- ³ Istituti Clinici Scientifici Maugeri IRCCS, Pulmonary Rehabilitation Unit of Telese Terme Institute, 82037 Telese Terme, Italy; silvestro.danna@icsmaugeri.it (S.E.D.); marco.mosella@icsmaugeri.it (M.M.)
- * Correspondence: pasquale.ambrosino@icsmaugeri.it (P.A.); mauro.maniscalco@icsmaugeri.it (M.M.)
- † These authors contributed equally to this work.
- ‡ These authors contributed equally to this work.

Table of contents:

Supplemental Figure 1	Flow chart of study participants.
-----------------------	-----------------------------------

Figure S1. Flow chart of study participants.



AMI: acute myocardial infarction; HF: heart failure; TIA: transient ischaemic attack; VTE: venous thromboembolism; PAD: peripheral artery disease; COPD: chronic obstructive pulmonary disease; CPET: cardiopulmonary exercise testing; FMD: flow-mediated dilation.