

Supplementary Figure S1




Figure S1. The expression levels of NRDC in different groups, including : **(A)** intraductal papillary mucinous neoplasm(IPMN) : n=6, chronic pancreatitis(CP) : n=23, pancreatic neuroendocrine tumor(PN) : n=2, Normal Control(NC):61, pancreatic ductal adenocarcinoma(PDAC) : n=112 . **(B)** Receiver operating characteristic curve (ROC) analyses of serum NRDC in TCGA and GTEx database. ns: none significance, ** $p < 0.01$, *** $p < 0.001$.

Supplementary Table S1

Table S1. Expression of NRDC in PDAC and its precursor lesions.

Diagnosis	<i>n</i>	Expression score mean ± SD	<i>P</i> value
intraductal papillary mucinous neoplasm(IPMN)	6	1308 ± 340	<i>p</i> =0.0054 (vs PDAC)
chronic pancreatitis(CP)	23	1132 ± 418	<i>p</i> =0.0006 (vs PDAC)
pancreatic neuroendocrine tumor(PNT)	2	747 ± 66	<i>p</i> =0.93965 (vs PDAC)
Normal Control(NC)	61	1483 ± 933	<i>p</i> <0.0001 (vs PDAC)
pancreatic ductal adenocarcinoma(PDAC)	112	771 ± 453	<i>p</i> <0.0001 (vs others)

SD, standard deviation; vs, versus