Supplemental Table S3. Trend in general anesthesia use from 2005 to 2020 according to insurance and clinical definitions

Fiscal year	N	Definition of general anesthesia, % of N (95%CI) General anesthesia claim code (L008) (insurance definition of general anesthesia)	L008 with NMBAs (clinical definition of general anesthesia)	L008 without NMBAs
2005	455	29.5 (25.3–33.6)	9.9 (7.2–12.6)	19.6 (15.9–23.2)
2006	550	29.3 (25.5–33.1)	7.5 (5.3–9.7)	21.8 (18.4–25.3)
2007	568	26.6 (23.0–30.2)	8.6 (6.3–10.9)	18.0 (14.8–21.1)
2008	744	23.5 (20.5–26.6)	5.1 (3.5–6.7)	18.4 (15.6–21.2)
2009	1109	21.3 (18.9–23.7)	5.2 (3.9–6.5)	16.1 (13.9–18.2)
2010	1863	22.5 (20.6–24.4)	4.7 (3.8–5.7)	17.8 (16.0–19.5)
2011	2567	23.8 (22.1–25.4)	5.9 (5.0–6.8)	17.8 (16.4–19.3)
2012	3494	21.8 (20.4–23.1)	4.4 (3.7–5.1)	17.3 (16.1–18.6)
2013	5274	16.7 (15.7–17.7)	4.3 (3.8–4.9)	12.4 (11.5–13.3)
2014	5707	13.9 (13.0–14.8)	4.4 (3.9–5.0)	9.5 (8.7–10.2)
2015	7264	12.9 (12.2–13.7)	3.6 (3.2–4.0)	9.3 (8.7–10.0)
2016	8955	12.4 (11.8–13.1)	3.5 (3.2–3.9)	8.9 (8.3–9.5)
2017	10511	12.2 (11.6–12.9)	3.2 (2.8–3.5)	9.1 (8.5–9.6)
2018	12527	12.5 (11.9–13.0)	3.7 (3.4–4.0)	8.8 (8.3–9.3)
2019	12109	12.1 (11.5–12.7)	3.6 (3.2–3.9)	8.6 (8.1–9.1)
2020	2483	11.5 (10.2–12.7)	3.3 (2.6–4.1)	8.1 (7.1–9.2)
total	76180	14.4 (14.2–14.7)	3.9 (3.8–4.1)	10.5 (10.3–10.7)

CI, confidence interval; NMBAs, neuromuscular blocking agents