

Supplemental Table S1. Comparison of substrate characteristics between patients with different periods on amiodarone before ablation

	Amiodarone > 6 months (N = 21)	Amiodarone more than 8 weeks and ≤ 6 months (N = 22)	<i>p</i> -value
RV endocardium			
Bipolar voltage*	1.8 ± 0.6	2.0 ± 0.7	0.222
Unipolar voltage*	4.9 ± 1.7	4.9 ± 1.2	0.999
Total activation time (ms)	167.4 ± 27.2	161.3 ± 34.2	0.496
Bipolar low voltage zone (cm ²)	40.4 ± 32.3	29.2 ± 20.9	0.185
Bipolar low voltage zone, %	17.3 ± 12.9	13.6 ± 8.0	0.262
Bipolar scar (cm ²)	19.5 ± 15.8	15.4 ± 11.2	0.332
Bipolar scar, %	8.8 ± 6.6	7.3 ± 4.3	0.385
Unipolar low voltage zone (cm ²)	65.2 ± 34.2	65.4 ± 21.2	0.977
Unipolar low voltage zone, %	26.6 ± 12.8	26.6 ± 7.9	0.999
Area with abnormal electrograms	17.7 ± 17.4	15.8 ± 13.7	0.692
RV epicardium	N=16	N=19	
Bipolar voltage*	1.5 ± 1.0	1.1 ± 0.4	0.152
Total activation time (ms)	212.3 ± 31.6	210.3 ± 21.0	0.825
Bipolar low voltage zone (cm ²)	97.6 ± 53.9	123.2 ± 56.0	0.180
Bipolar low voltage zone, %	33.1 ± 22.3	38.9 ± 18.8	0.410
Bipolar scar (cm ²)	51.1 ± 34.0	61.2 ± 34.6	0.390
Bipolar scar, %	17.4 ± 13.4	18.6 ± 9.7	0.756

Area with abnormal electrograms	24.7 ± 30.1	31.8 ± 28.1	0.434
------------------------------------	-------------	-------------	-------

*The average of bipolar or unipolar median voltage.

ARVC, arrhythmogenic right ventricular cardiomyopathy; RV, right ventricular; VT, ventricular tachycardia

“