

Table S1. Comparative analysis of the ESSDAI domains between the two groups.

	PN+ Group	PN- Group	<i>p</i> -Value
	No = 31 (25.61%)	No = 90 (74.38%)	
Constitutional, <i>n</i> (%)	0	0	-
Lymphadenopathy, <i>n</i> (%)	5 (16.12%)	20 (22.22%)	0.328 ^a
Glandular, <i>n</i> (%)	15 (48.38%)	68 (75.55%)	0.006 ^a
Articular, <i>n</i> (%)	19 (61.29%)	55 (61.11%)	0.581 ^a
Cutaneous, <i>n</i> (%)	12 (38.70%)	22 (24.44%)	0.471 ^a
Pulmonary, <i>n</i> (%)	10 (32.25%)	17 (18.88%)	0.138 ^a
Renal, <i>n</i> (%)	10 (32.25%)	13 (14.44%)	0.031 ^a
Muscular, <i>n</i> (%)	4 (12.90%)	15 (16.66%)	0.429 ^a
Hematological, <i>n</i> (%)	13 (41.93%)	38 (42.22%)	0.575 ^a
Biological, <i>n</i> (%)	13 (41.93%)	19 (21.11%)	0.023 ^a

Abbreviations: PN, peripheral neuropathy. ^a Chi- Square; statistical significance <0.05.