

Supplementary Part (Subgroups):

ST1. Prevalence of P4 Risk Categories and Correlation with Other Scale Items and Scores (subgroups)

Specialized care

	Total (n=119)		Women (n=71)		Men (n=48)		Intercorrelations (convergent validity) **							
	no risk	risk	no risk	risk	no risk	risk	SBQ-R_1 (lifetime SI/attempts)	SBQ-R_2 (12-month SI)	SBQ-R_3 (threatened suicide)	SBQ-R_4 (suicide likelihood)	SBQ-R _sum	PMH _sum	PHQ9 _sum	PHQ9_ 09
P4_1 (past attempt)	84 (70.6%)	35 (29.4%)	46 (64.8%)	25 (35.2%)	38 (79.2%)	10 (20.8%)	.756**	.284**	.316**	.318**	.371**	.006	.142	.284**
P4_2 (suicide plan)	43 (36.1%)	76 (63.9%)	22 (31.0%)	49 (69%)	21 (43.8%)	27(56.3%)	.521**	.603**	.504**	.319**	.609**	.076	.060	.246**
P4_3 (suicide likelihood)	102 (85.7%)	17 (14.3%)	59 (83.1%)	12 (16.9%)	43 (89.6%)	5 (10.4%)	.089	.201*	.203*	.374**	.133	-.131	.204*	.310**
P4_4 (preventive factors)	105 (88.2%)	14 (11.8%)	63 (88.7%)	8 (11.3%)	42 (87.5%)	6 (12.5%)	.019	.064	.086	.112	.082	-.205*	.030	.132
P4_risk (minimal, lower, higher)	34 (28.6%)	lower: 61 (51.3%) higher: 24 (20.2%)	17 (23.9%)	lower: 39 (54.9%) higher: 15 (21.2%)	17 (35.4%)	lower: 22 (45.8%) higher: 9 (18.8%)	.335**	.405**	.345**	.420**	.448**	-.121	.124	.365**

** correlation is significant at P < .01 level, *corelation is significant at p< .05

Primary care

	Total (n=104)		Women (n=68)		Men (n=36)		Intercorrelations (convergent validity) *							
	no risk	risk	no risk	risk	no risk	risk	SBQ-R_1 (lifetime SI/attempts)	SBQ-R_2 (12-month SI)	SBQ-R_3 (threatened suicide)	SBQ-R_4 (suicide likelihood)	SBQ-R _sum	PMH _sum	PHQ9 _sum	PHQ9_ 09
P4_1 (past attempt)	76 (73.1%)	28 (26.9%)	50 (73.5%)	18 (26.5%)	26 (72.2%)	10 (27.8%)	.725**	.291**	.295**	.228*	.309**	.06	.07	.18
P4_2 (suicide plan)	41 (39.4%)	63 (60.6%)	24 (35.5%)	44 (64.7%)	17 (47.2%)	19 (52.8%)	.563**	.505**	.448**	.185	.388**	-.10	.212*	.216*
P4_3 (suicide likelihood)	98 (94.2%)	6 (5.7%)	65 (95.6%)	3 (3.4%)	33 (91.7%)	3 (8.4%)	.252**	.293**	.231*	.400**	.196*	-.05	.203*	.422**
P4_4 (preventive factors)	96 (93.2%)	7 (6.8%)	64 (95.5%)	3 (4.5%)	32 (88.9%)	4 (11.1%)	.049	.010	.061	-.048	-.003	-.053	.053	.129
P4_risk (minimal, lower, higher)	34 (33.0%)	lower: 58 (56.3%) higher: 11 (10.7%)	19 (28.4%)	lower: 42 (62.7%) higher: 6 (9%)	15 (34.7%)	lower: 16 (44.4%) higher: 5 (13.9%)	.600**	.492**	.408**	.229*	.354**	-.10	.283**	.358**

** correlation is significant at $P < .01$ level, *corelation is significant at $p < .05$

P4_1: Have you ever attempted to harm yourself in the past? (no/yes)

P4_2: Have you thought about how you might actually hurt yourself? (no/yes)

P4_3: There's a big difference between having a thought and acting on a thought. How likely do you think it is that you will act on these thoughts about hurting yourself or ending your life some time over the next month? (not at all likely/somewhat likely/ very likely)

P4_4: Is there anything that would prevent or keep you from harming yourself? (no/yes)

ST2. Qualitative evaluation (Subgroups) of Suicidal Ideation Plans and Preventive Factors

<u>Specialized setting</u>			<u>Primary care</u>		
Suicidal Ideations Plans	N	%	Suicidal Ideations Plans	N	%
medication overdose	27	29.7	medication overdose	20	27.8
cutting oneself	18	19.8	intentional vehicular accident	14	19.4
intentional vehicular accident	15	16.5	cutting oneself	11	15.3
others	13	14.3	others	8	11.1
hanging	6	6.6	falls	8	11.1
falls	6	6.6	no answer	7	9.7
no answer	5	5.5	hanging	3	4.2
using a gun	1	1.1	using a gun	1	1.4
Total	91	100	Total	72	100

<u>Specialized setting</u>			<u>Primary care</u>		
Preventive Factors	N	%	Preventive Factors	N	%
family	78	66.1	family	69	69.0
future hope	20	16.9	future hope	13	13.0
faith	6	5.1	faith	6	6.0
fear of failiing	4	3.4	fear of failiing	4	4.0
others	10	8.5	others	8	8.0
Total	118	100	Total	100	100