Table S1.S Questionnaire

Gender Age Zip Code (living) Zip Code (working)	male female	
Working status	Private dentist Private/NHS NSH	
From the start of the COVID-19 you had	No symptoms You resulted COVID-19 positive You were hospitalized for COVID I had one/more symptoms	Fever Cough Fatigue Short Breath Nasal congestion Headache Rhinorrhea Sore throat Diffuse pain Diarrhea Anosmia Ageusia Conjunctivitis
Only if you work in the NHS, are you currently working?	Yes No	
From the 21st February	You kept working as usual You limited your activity to emergencies You have stopped all activities	
If you have limited the activity to emergencies, when did you start limiting?	Between February 21-23 Between 24 February and 1 March Between March 2-6 Between March 7-14 After March the 14th	
If you have stopped the activity, when it is happened?	Between February 21-23 Between 24 February and 1 March Between March 2-6 Between March 7-14 After March the 14th	
In the case that you have continued working after February 21st, which of the following precautionary measures have you taken?	None Phone Triage Appointments reduced so as not to saturate the waiting room Postponement of therapies in elderly patients, or with systemic diseases	

	Handle disinfection several times a day Disinfection of push buttons, POS, chairs, several times a day Verify the patient's current health status on access Detecting the patient's body temperature Detection of body temperature of all co-workers and leave of those with a temperature above 37.5°. Washing the patient's hands Space of at least one meter between patients Mask for the patient Frequent ventilation of waiting rooms Removal of magazines and books from the waiting area Storage of coats, bags and other items outside the operating area Pre-operative rinse with mouthwash containing 1% hydrogen peroxide Pre-operative rinse with mouthwash containing chlorhexidine 0.12-0.2% Pre-operative rinse with mouthwash containing 0.2-1% iodopovidone Pre-operative rinse with mouthwash containing alcohol and essential oils Pre-operative rinse with mouthwash containing Cetylpyridinium chloride at 0.05-0.10% Rinse with diluted mouthwash Ventilation of the operating area for at least 10 minutes after each patient Surface disinfection with 70% ethyl alcohol Disinfection of surfaces with 0.5% sodium hypochlorite Usual disinfectant with others active ingredients Washing operators' hands before and after each procedure Removal of all disposable protective devices and disinfection of non-disposable devices
Which of the following protections have you adopted?	Surgical mask FFP2 or FFP3 facial filters Disposable headset Sterile microfiber disposable gown Water-repellent TNT disposable gown Disposable gown Safety glasses or visor Sterile disposable gloves Disposable gloves Rotating instrument with anti-retraction valve
Did you follow a course on Covid-19?	Yes No
Do you think that you know enough on COVID?	Yes No
	Unlikely Very unlikely

Do you believe that the infection by SARS-CoV-2 is a risk for the dentist?	Likely Very likely
Do you believe that the infection by SARS-CoV-2 is a risk for the dentist?	Unlikely Very unlikely Likely Very likely
How sure are you that you can avoid becoming infected with SARS-CoV-2 during work?	No confident Enough confident A bit confident Confident
Do you believe that in a health emergency situation such as the current one, the risk of infection transmission in the dental practice is	Less than the risk runs going in a supermarket Comparable to the risk runs going in a supermarket Higher than the risk runs going in a supermarket