

Supplementary Table S1.*Presence of gastrointestinal disorders in the total sample.*

Gastrointestinal disorders	<i>n</i> (%)
Infectious diarrhea	287 (23.0)
Nonspecific abdominal pain	246 (19.7)
Dyspepsia	132 (10.6)
Gastroesophageal reflux	138 (11.1)
Significant Flatulence	67 (5.4)
Irritable Bowel Syndrome	46 (3.7)
Dyschezia	18 (1.4)
Inflammatory Bowel Disease	13 (1.0)
Celiac Disease	11 (0.9)
Ulcerative Colitis	9 (.7)
Peptic Ulcer Disease	5 (.4)
Crohn's disease	4 (.3)

Supplementary Table S2.*Percentiles of the GSSS.*

	Percentile	Severity level of gastrointestinal symptoms	Factor 1	Factor 2	Total SSGS
Total Sample	29	Very Low	0	0	1
	49	Low	1	0	2
	69	Medium-low	2	1	3
	79	Medium-High	3	1	4
	89	High	4	2	5
	99	Very High	7	6	12
M (SD) Total Sample			1.63 (1.74)	.73 (1.21)	2.37 (2.54)
Females	29	Very Low	0	0	1
	49	Low	1	0	2
	69	Medium-Low	2	1	3
	79	Medium-High	3	1	4
	89	High	4	2	6
	99	Very High	8	6	12
M (SD) Females			1.79 (1.82)	.76 (1.24)	2.55 (2.64)
Males	29	Very Low	0	0	0
	49	Low	1	0	1
	69	Medium-Low	2	1	3
	79	Medium-High	2	1	3
	89	High	3	2	4
	99	Very High	6	6	10
M (SD) Males			1.18 (1.38)	.66 (1.09)	1.84 (2.13)
Contrast Statistics between sex (Effect Size)			P value		Hedges' g
Total SSGS Females vs. Total SSGS Males			$p < .05$.28
Factor 1 SSGS Females vs. Factor 1 SSGS Males			$p < .05$.35
Factor 2 SSGS Females vs. Factor 2 SSGS Males			$p > .05$.08

Notes. M=Mean; SD= Standard Deviation.