Supplementary Table S1. Post-operative 30-day outcomes among patients with and without SCI evolution. Footnotes: ICU: intensive care unit; SCI: spinal cord ischemia

Post-operative outcomes (n, %)	SCI group (23 patients)	Non-SCI group (23 patients)	P
Mortality	2 (8.7%)	0 (0.0%)	0.14
SCI	23 (100%)	0 (0.0%)	-
-Grade 1	4 (17.4%)	-	-
-Grade 2	14 (60.9%)	-	-
-Grade 3	5 (21.7%)	-	-
-Late evolution	7 (30.4%)	-	-
-Recovery	19 (82.6%)	-	-
-Partial	12 (52.2%)	-	-
-Total	7 (30.4%)	-	-
Stroke	0 (0.0%)	0 (0.0%)	-
Mesenteric ischemia	2 (8.7%)	0 (0.0%)	0.14
Myocardial infarction	0 (0.0%)	0 (0.0%)	-
Acute kidney injury	10 (43.5%)	3 (13.0%)	0.02
Bleeding	6 (26.1%)	1 (4.3%)	0.04
-Needing reintervention	3 (13.0%)	0 (0.0%)	0.07
ICU length (days)	11 (IQR 7, range 2-30)	3 (IQR 2, range 1-13)	<0.001
Length of hospital stay (days)	17 (IQR 20, range 2-65)	13 (IQR 8, range 5-43)	0.02