

Sex Differences in Prognostic Markers: Exploring Outcome Variability after Mechanical Thrombectomy in Large Vessel Occlusion Stroke

Hannes Schacht ^{1,*}, Alexander Neumann ¹, Nora Petersen ², Lis Merrit Ehm ², Maria Marburg ², Christine Matthis ³, Ulf Jensen-Kondering ¹, Peter Schramm ¹, Jens Minnerup ², Georg Royl ² and Philipp J. Koch ^{2,4}

¹ Department of Neuroradiology, University Hospital Schleswig-Holstein, Campus Lübeck, 23538 Lübeck, Germany; Alexander.Neumann@uksh.de (A.N.); Ulf.Jensen-Kondering@uksh.de (U.J.-K.); Peter.Schramm@uksh.de (P.S.)


² Department of Neurology, University Hospital Schleswig-Holstein, Campus Lübeck, 23538 Lübeck, Germany; nora.petersen@student.uni-luebeck.de (N.P.); LisMerrit.Ehm@uksh.de (L.M.E.); MariaJoana.Marburg2@uksh.de (M.M.); Jens.Minnerup@uksh.de (J.M.); Georg.Royl@uksh.de (G.R.)

³ Department of Social Medicine and Epidemiology, University Hospital Schleswig-Holstein, Campus Lübeck, 23538 Lübeck, Germany; Christine.Matthis@uksh.de (C.M.)

⁴ Department of Neurology and Experimental Neurology, Charité - University Medicine Berlin, corporate member of Freie Universität Berlin and Humboldt-Universität zu Berlin, 10117 Berlin, Germany; philipp-johannes.koch@charite.de (P.J.K.)

* Correspondence: Hannes.Schacht@uksh.de; Tel.: +49-45150076183

a Score on the Modified Rankin Scale at discharge



b Score on the Modified Rankin Scale at 90 days




Figure S1. Bar graphs illustrate the overall differences between female and male patients in the modified Rankin Scale (mRS) at discharge (a) and 90 days following stroke (b).