Supplementary Table S2 Baseline comparison between participants with EEG recordings and those without EEG recordings within the full randomized cohort.

Characteristic	With EEG recordings (n=56)	Without EEG recordings (n=53)	P Value
Age (years)	57.0 ± 4.7	57.7 ± 4.5	0.430
Sex, n (%)			0.614
Female	27	23	
Male	29	30	
BMI (kg/m^2)	21.9 ± 1.1	21.8 ± 1.3	0.571
Procedure			0.849
Single	49	47	
Bilobectomy	7	6	
Duration of surgery (min)	108.4 ± 16.5	108.8 ± 14.5	0.869
Dose of propofol (mg)	966.6 ± 67.7	951.4 ± 50.2	0.187
Dose of remifentanil (mg)	1.3 ± 0.3	1.2 ± 0.2	0.668
Histological subtype, n (%)			0.692
Adenocarcinoma	44	43	
Squamous-cell carcinoma	11	8	
Others [a]	1	2	
Tumour maximum diameter (cm), mean ± SD	3.2 ± 0.5	3.1 ± 0.5	0.220
Lymph-node-dissection pattern, n (%)			0.407
Systematic mediastinal lymph-node dissection	36	38	
Lymph-node sampling	20	15	

Data are expressed as number (%) or mean ± standard deviation. BMI, body mass index; TEAS, transcutaneous electrical acupoint stimulation.

[a] include adenosquamous carcinoma and large-cell carcinoma.